

---

# **Twinrix**

**Version GDSv17-IPIv11**

# Twinrix

## Inactivated hepatitis A and rDNA hepatitis B vaccine (adsorbed) Suspension for injection

### Qualitative and quantitative composition

Suspension for injection.

One dose (1.0 ml) contains:

Hepatitis A virus (inactivated) <sup>1,2</sup>	720 ELISA Units
Hepatitis B surface antigen <sup>3,4</sup>	20 micrograms
<sup>1</sup> Produced on human diploid (MRC-5) cells	
<sup>2</sup> Adsorbed on aluminium hydroxide, hydrated	0.05 milligrams Al <sup>3+</sup>
<sup>3</sup> Produced in yeast cells ( <i>Saccharomyces cerevisiae</i> ) by recombinant DNA technology	
<sup>4</sup> Adsorbed on aluminium phosphate	0.4 milligrams Al <sup>3+</sup>

Twinrix is a white, slightly milky liquid presented in a glass pre-filled syringe

### CLINICAL INFORMATION

#### Indications

Twinrix is indicated for use in non-immune adults, adolescents and children aged from 1 year upwards, who are at risk of both hepatitis A and hepatitis B infection.

#### Dosage and Administration

##### Dosage

A dose of 1.0 ml Twinrix is recommended for adults, adolescents and children aged from 1 year upwards.

##### Primary vaccination schedules

- *Adults and adolescents of 16 years of age and above*

The standard primary course of vaccination with Twinrix consists of three doses, the first administered at the elected date, the second one month later and the third six months after the first dose.

In exceptional circumstances in adults, when travel is anticipated within one month or more after initiating the vaccination course, but where insufficient time is available to allow the standard 0, 1, 6 month schedule to be completed, a schedule of three intramuscular injections given at 0, 7 and 21 days may be used. When this schedule is applied, a fourth dose is recommended 12 months after the first dose.

- *Children and adolescents of 1 to 15 years of age*

The standard primary course of vaccination with Twinrix consists of two doses, the first is administered at the elected date and the second between six and twelve months after the first dose. As protection against hepatitis B infection will not be obtained in all vaccinees until after the second dose, it is important that the second dose be administered to assure protection against hepatitis B infection.

The recommended schedule should be adhered to. Once initiated, the primary course of vaccination should be completed with the same vaccine.

##### Booster dose

Long-term antibody persistence data following vaccination with Twinrix in adults and adolescents (0, 1, 6 month) are available up to 20 years after vaccination.

Long-term antibody persistence data following vaccination with Twinrix (0, 6 month) are available up to 10 years in individuals aged 1-11 years at primary

vaccination and up to 15 years in individuals aged 12-15 years at primary vaccination (see section *Pharmacodynamics*).

The anti-HBs and anti-HAV antibody titres observed following a primary vaccination course with the combined vaccine are in the range of what is seen following vaccination with the monovalent vaccines. General guidelines for booster vaccination can therefore be drawn from experience with the monovalent vaccines.

##### *Hepatitis B*

The need for a booster dose of hepatitis B vaccine in healthy individuals who have received a full primary vaccination course has not been established; however some official vaccination programmes currently include a recommendation for a booster dose of hepatitis B vaccine and these should be respected.

For some categories of subjects or patients exposed to HBV (e.g. haemodialysis or immunocompromised patients) a precautionary attitude should be considered to ensure a protective antibody level  $\geq 10$  IU/l.

##### *Hepatitis A*

It is not yet fully established whether immunocompetent individuals who have responded to hepatitis A vaccination will require booster doses, as protection in the absence of detectable antibodies may be ensured by immunological memory. Guidelines for boosting are based on the assumption that antibodies are required for protection.

In situations where a booster dose of both hepatitis A and hepatitis B are desired, Twinrix can be given. Alternatively, subjects primed with Twinrix may be administered a booster dose of either of the monovalent vaccines. The safety and immunogenicity of Twinrix administered as a booster dose following a two dose primary vaccination course have not been evaluated.

##### **Method of administration**

Twinrix should be injected intramuscularly into the deltoid region of the upper arm in adults, adolescents and children. The anterolateral thigh may be used in infants.

Since intradermal injection or intramuscular administration into the gluteal muscle could lead to a suboptimal response to the vaccine, these routes should be avoided. Exceptionally, Twinrix can be administered subcutaneously to subjects with thrombocytopenia or bleeding disorders since bleeding may occur following an intramuscular administration to these subjects. However, this route of administration may result in suboptimal immune response to the vaccine.

##### **Contraindications**

Twinrix should not be administered to subjects with known hypersensitivity to any constituent of the vaccine, or to subjects having shown signs of hypersensitivity after previous administration of Twinrix or the monovalent hepatitis A or hepatitis B vaccines.

##### **Warnings and Precautions**

As with other vaccines, the administration of Twinrix should be postponed in subjects suffering from acute severe febrile illness.

Syncope (fainting) can occur following, or even before, any vaccination as a psychogenic response to the needle injection. It is important that procedures are in place to avoid injury from faints.

It is possible that subjects may be in the incubation period of a hepatitis A or hepatitis B infection at the time of vaccination. It is not known whether Twinrix will prevent hepatitis A and hepatitis B in such cases.

The vaccine will not prevent infection caused by other agents such as hepatitis C and hepatitis E and other pathogens known to infect the liver.

Twinrix is not recommended for postexposure prophylaxis (e.g. needle stick injury).

The vaccine has not been tested in patients with impaired immunity. In haemodialysis patients and persons with an impaired immune system, adequate anti-HAV and anti-HBs antibody titres may not be obtained after the primary

immunisation course and such patients may therefore require administration of additional doses of vaccine.

As with all injectable vaccines, appropriate medical treatment and supervision should always be readily available in case of a rare anaphylactic event following the administration of the vaccine.

Twinrix should under no circumstances be administered intravascularly.

## Interactions

No data on concomitant administration of Twinrix with specific hepatitis A immunoglobulin or hepatitis B immunoglobulin have been generated. However, when the monovalent hepatitis A and hepatitis B vaccines were administered concomitantly with specific immunoglobulins, no influence on seroconversion was observed although it may result in lower antibody titres.

Clinical studies have demonstrated that Twinrix can be administered concomitantly with diphtheria, tetanus, acellular pertussis, inactivated poliomyelitis, *Haemophilus influenzae* type b (DTPa-IPV/Hib) or Measles-Mumps-Rubella vaccines in the second year of life. In these trials, the injectable vaccines were given at different injection sites.

Although the concomitant administration of Twinrix and other vaccines has not specifically been studied, it is anticipated that, if different syringes and other injection sites are used, no interaction will be observed.

It may be expected that in patients receiving immunosuppressive treatment or patients with immunodeficiency, an adequate response may not be achieved.

## Pregnancy and Lactation

### Pregnancy

Twinrix should be used during pregnancy only when clearly needed and when the possible advantages outweigh the possible risks for the foetus.

The effect of Twinrix on embryo-foetal, peri-natal and post-natal survival and development has not been prospectively evaluated in clinical trials.

The effect of Twinrix on embryo-foetal, peri-natal and post-natal survival and development has been assessed in rats. Such animal studies do not indicate direct or indirect harmful effects with respect to fertility, pregnancy, embryonal/foetal development, parturition or post-natal development.

### Lactation

Adequate human data on use during lactation and adequate animal reproduction studies are not available. Twinrix should therefore be used with caution in breastfeeding women.

## Effects on Ability to Drive and Use Machines

The vaccine is unlikely to produce an effect on the ability to drive and use machines.

## Adverse Reactions

Local and general adverse events reported following primary vaccination with Twinrix were categorised by frequency.

Adverse reactions reported are listed according to the following frequency:

Very common:	≥ 1/10
Common:	≥ 1/100 to < 1/10
Uncommon:	≥ 1/1000 to < 1/100
Rare:	≥ 1/10000 to < 1/1000
Very rare:	< 1/10000

- Adults and adolescents of 16 years of age and above

The safety profile presented below is based on data from more than 6,000 subjects who received either the standard 0, 1, 6 month schedule or the accelerated 0, 7, 21 days schedule.

Clinical trials		
System Organ Class	Frequency	Adverse reactions
Infections and infestations	Uncommon	Upper respiratory tract infection
Blood and lymphatic system disorders	Rare	Lymphadenopathy
Metabolism and nutrition disorders	Rare	Decreased appetite
Nervous system disorders	Very common	Headache
	Uncommon	Dizziness
	Rare	Hypoaesthesia, paraesthesia
Vascular disorders	Rare	Hypotension
Gastrointestinal disorders	Common	Gastrointestinal symptoms (such as diarrhoea, nausea, vomiting)
Skin and subcutaneous tissue disorders	Rare	Rash, pruritus
	Very rare	Urticaria
Musculoskeletal and connective tissue disorders	Uncommon	Myalgia
	Rare	Arthralgia
General disorders and administration site conditions	Very common	Pain and redness at the injection site, fatigue
	Common	Swelling at the injection site, injection site reaction, malaise
	Uncommon	Fever (≥ 37.5°C)
	Rare	Influenza like illness, chills
Post-marketing data*		
System Organ Class	Adverse reactions	
Infections and infestations	Meningitis	
Blood and lymphatic system disorders	Thrombocytopenia, thrombocytopenic purpura	
Immune system disorders	Anaphylaxis, allergic reactions including anaphylactoid reactions and mimicking serum sickness	
Nervous system disorders	Encephalitis, encephalopathy, neuritis, neuropathy, paralysis, convulsions	
Vascular disorders	Vasculitis	
Skin and subcutaneous tissue disorders	Angioneurotic oedema, lichen planus, erythema multiforme	
Musculoskeletal and connective tissue disorders	Arthritis, muscular weakness	
General disorders and administration site conditions	Immediate injection site pain, stinging and burning sensation	

\* These adverse reactions have been reported with either Twinrix (given as a 0, 1, 6-month schedule or a 0, 7, 21 days schedule) or with GlaxoSmithKline monovalent hepatitis A or B vaccines.

In a comparative study it was noted that the frequency of solicited adverse events following the administration of Twinrix is not different from the frequency of solicited adverse events following the administration of the monovalent vaccines.

In a clinical trial where Twinrix was administered at 0, 7, 21 days, solicited general symptoms were reported with the same categories of frequency as defined

above. After a fourth dose given at month 12, the incidence of systemic adverse reactions was comparable to that seen after vaccination at 0, 7, 21 days.

- *Children and adolescents of 1 to 15 years of age inclusive*

Clinical trials involved the administration of 1537 doses of Twinrix as a two dose schedule to 778 subjects from 1 year up to and including 15 years of age.

Clinical trials		
System Organ Class	Frequency	Adverse reactions
Metabolism and nutrition disorders	Very common	Appetite lost
Psychiatric disorders	Very common	Irritability
Nervous system disorders	Very common	Headache
	Common	Drowsiness
Gastrointestinal disorders	Common	Gastrointestinal symptoms
General disorders and administration site conditions	Very common	Fatigue, pain and redness at the injection site
	Common	Fever, swelling at the injection site
Post-marketing data*		
System Organ Class	Frequency	Adverse reactions
Immune system disorders	Very rare	Allergic reactions including anaphylactic and anaphylactoid reactions
Nervous system disorders	Very rare	Syncope or vasovagal responses to injection, hypoesthesia

\* These adverse reactions were reported during the post-marketing surveillance following vaccination with Twinrix given as a 2-dose schedule.

## Overdose

Cases of overdose have been reported during post-marketing surveillance after administration of Twinrix. Adverse reactions reported following overdosage were similar to those reported with normal vaccine administration.

## PHARMACOLOGICAL PROPERTIES

### Pharmacodynamics

Pharmaco-therapeutic group: Hepatitis vaccines, ATC code J07BC20.

Twinrix confers immunity against HAV and HBV infection by inducing specific anti-HAV and anti-HBs antibodies.

- *Children and adolescents of 1 to 15 years of age*

In clinical studies involving subjects from 1 to 15 years old, seropositivity rates for anti-HAV antibodies were 99.1% one month after the first dose and 100% after the second dose given at month 6 (i.e. month 7). Seropositivity rates for anti-HBs antibodies were 74.2% one month after the first dose and 100% after the second dose given at month 6 (i.e. month 7). The anti-HBs seroprotection rates (titres  $\geq$  10 mIU/ml) at these time points were 37.4% and 98.2% respectively.

In a comparative study conducted in subjects (from 12 years up to and including 15 years of age) vs an alternative schedule of 3 doses with the combined vaccine containing 360 ELISA Units inactivated HA virus and 10  $\mu$ g HBsAg in a dose volume of 0.5 ml, seroprotection rates for anti-HBs at intermediate time points before the 2<sup>nd</sup> dose of Twinrix were lower compared to those obtained with the alternative schedule comprising 3 doses, but non-inferiority was shown after completion of the schedule (month 7).

Anti-HAV and anti-HBs antibodies have been shown to persist for at least 10 years following the initiation of a 0, 6 month schedule of Twinrix. After 10 years, the

anti-HAV seropositivity rates and the anti-HBs seroprotection rates in subjects aged 1-11 years at primary vaccination were 100% and 77.3% respectively. After 15 years, the anti-HAV seropositivity rates and the anti-HBs seroprotection rates in subjects aged 12-15 years at primary vaccination, were 100% and 81.1% respectively. A challenge dose of a HBV vaccine was given to a limited number of subjects (n=8) whose anti-HBs antibody concentrations decreased to < 10 mIU/ml and all mounted an anamnestic response.

After 15 years, the immune response for both antigen components was comparable to that seen after a 3-dose regimen of the combined vaccine containing 360 ELISA Units of inactivated hepatitis A virus and 10  $\mu$ g of recombinant hepatitis B surface antigen in a dose volume of 0.5 ml.

In a clinical study conducted among 12 years up to and including 15 years old subjects who received the second dose of Twinrix at month 12, seropositivity rates for anti-HAV were 99.0% and seropositivity rates for anti-HBs were 99.0% at month 13 with seroprotection rates of 97.0%.

In a 6 year long-term follow-up study involving subjects aged 12-15 years at primary vaccination, anti-HAV seropositivity rates were 100% following a 0, 6 month or a 0, 12 month schedule. The anti-HBs seroprotection rates were 84.8% and 92.9%, respectively.

- *Adults and adolescents of 16 years of age and above*

In subjects aged 16 years and above administered a 3-dose schedule of Twinrix, protection against hepatitis A and hepatitis B develops within 2-4 weeks. In the clinical studies, specific humoral antibodies against hepatitis A were observed in approximately 94% of the adults one month after the first dose and in 100% one month after the third dose (i.e. month 7). Specific humoral antibodies against hepatitis B were observed in 70% of the adults after the first dose and approximately 99% after the third dose.

For use in exceptional circumstances in adults, the 0, 7 and 21 day primary schedule plus a fourth dose at month 12 results in 82% and 85% of vaccinees having seroprotective levels of anti-HBV antibodies at 1 and 5 weeks respectively following the third dose. One month after the fourth dose, all vaccinees demonstrated seroprotective levels of antibody. Seropositivity rates for anti-HAV antibodies were 100% and 99.5% at 1 and 5 weeks respectively following the third dose, and reached 100% one month after the fourth dose.

In a clinical study conducted in subjects over 40 years of age, the seropositivity rate for anti-HAV antibodies and seroprotection rate against hepatitis B following Twinrix on a 0, 1, 6 month schedule were compared with the seropositivity and seroprotection rates of monovalent hepatitis A and B vaccines when administered separately.

The seroprotection rates against hepatitis B after the administration of Twinrix were 92% and 57% at 7 and 48 months following the first dose respectively, versus 80% and 40% after the GlaxoSmithKline Biologicals monovalent 20  $\mu$ g hepatitis B vaccine, and 71% and 27% after another licensed monovalent 10  $\mu$ g hepatitis B vaccine. In all groups, anti-HBs antibody concentrations decreased as age and body mass index increased; concentrations were also lower in males compared with females.

The seropositivity rates for anti-HAV antibodies after Twinrix were 97% at both 7 and 48 months following the first dose versus 99% and 94% after the GlaxoSmithKline Biologicals monovalent hepatitis A vaccine and 99% and 96% after another licensed monovalent hepatitis A vaccine.

Subjects received an additional dose of Twinrix to assess the immune memory 48 months after the first dose of the primary vaccination course with the same vaccine. One month after this dose, 95% of subjects elicited anti-HBV antibody concentration  $\geq$  10 mIU/ml and Geometric Mean Concentrations (GMC) increased by 179-fold (GMC of 7233.7 mIU/ml) indicative of an immune memory response.

In two long-term clinical studies conducted in adults, 20 years after the primary vaccination with Twinrix the anti-HAV seropositivity rates were 100% and 96% respectively and the anti-HBs seroprotection rates were 94% and 92%, respectively (n=43).

Non-Clinical Information Pre-clinical data reveal no special hazard for humans based on general safety studies (see section *Pregnancy and Lactation*).

## PHARMACEUTICAL INFORMATION

### List of Excipients

Sodium chloride and water for injections. Aminoacids for injection, formaldehyde, neomycin sulphate and polysorbate 20 are present as residual from the manufacturing process.

### Shelf Life

The expiry date is indicated on the label and packaging.

### Storage

Twinrix should be stored at +2 °C to +8 °C.

Store in the original package in order to protect from light.

**Do not freeze;** discard if the vaccine has been frozen.

The storage conditions are detailed on the packaging

### Nature and Contents of Container

1 ml of suspension in a pre-filled syringe (type I glass) with a plunger stopper (butyl rubber) and with a rubber tip cap.

The tip cap and rubber plunger stopper of the pre-filled syringe are not made with natural rubber latex.

Not all presentations are available in every country.

### Incompatibilities

Twinrix should not be mixed with other vaccines in the same syringe.

### Use and Handling

The vaccine should be re-suspended before use. When re-suspended, the vaccine will have a uniform hazy white appearance.

Upon storage, a fine white deposit with a clear colourless layer above may be observed.


### Re-suspension of the vaccine to obtain a uniform hazy white suspension

The vaccine can be re-suspended following the steps below.

1. Hold the syringe upright in a closed hand.
2. Shake the syringe by tipping it upside down and back again.
3. Repeat this action vigorously for at least 15 seconds.
4. Inspect the vaccine again:
  - a. If the vaccine appears as a uniform hazy white suspension, it is ready to use – the appearance should not be clear.
  - b. If the vaccine still does not appear as a uniform hazy white suspension - tip upside down and back again for at least another 15 seconds - then inspect again.


The vaccine should be inspected visually for any foreign particulate matter and/or abnormal physical appearance prior to administration. In the event of either being observed, do not administer the vaccine.

[Instructions for the pre-filled syringe](#)



Hold the syringe by the barrel, not by the plunger.

Unscrew the syringe cap by twisting it anticlockwise.



To attach the needle, connect the hub to the Luer Lock Adaptor and rotate a quarter turn clockwise until you feel it lock.

Do not pull the syringe plunger out of the barrel. If it happens, do not administer the vaccine.

### Disposal

Any unused product or waste material should be disposed of in accordance with local requirements.

**Version number: GDS17/IP11**

**Date of issue: 05-May-2023**