

e-Banner	Project: CO-0051986	Document: PPC-0029975	Version: 3
	Site Code: 6200000081704	Operator: SRM58890	Date/Time Created: 12.Sep.2022 17:45 GMT+1

Pharma code Ref. No. 1173 Pharma code Ref. No. 1173



- breast enlargement or tenderness (*gynecomastia*)
- dizziness when taken with tamsulosin.

Uncommon side effects

- These may affect up to 1 in 100 men taking Avodart:*
- heart failure (heart becomes less efficient at pumping blood around the body. You may have symptoms such as shortness of breath, extreme tiredness and swelling in your ankles and legs)
 - hair loss (usually from the body) or hair growth.

Not known side effects

- The frequency cannot be estimated from the available data:
- depression
 - pain and swelling in your testicles.

5 How to store Avodart

- Keep this medicine out of the sight and reach of children.
- Don't store Avodart above 30°C.
- Do not use this medicine after the expiry date which is stated on the carton or the foil blister strip. The expiry date refers to the last day of that month.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6 Contents of the pack and other information

What Avodart contains

- The active substance is dutasteride.** Each soft capsule contains 0.5 mg dutasteride. The other ingredients are:
- inside the capsule: mono-di-glycerides of caprylic/capric acid and butylated hydroxytoluene (E321).
 - capsule shell: gelatin, glycerol, titanium dioxide (E171), iron oxide yellow (E172), triglycerides (medium chain), lecithin (may contain soya oil).

What Avodart looks like and contents of the pack
Avodart soft capsules are oblong opaque, yellow, soft gelatin capsules marked with GX CE2. They are available in packs of 10, 30, 50, 60 and 90 capsules. Not all pack sizes may be available.

Marketing Authorisation Holder:
GlaxoSmithKline UK Limited
980 Great West Road, Brentford, Middlesex, United Kingdom

Manufacturer:

Delpharm Poznan Spolka Akcyjna, Poznan, Poland
Trade marks are owned by or licensed to the GSK group of companies.

© 2022 GSK group of companies or its licensor.
To report Product Complaints or Adverse Event/s associated with the use of GSK products, please contact us via: gulf.safety@gsk.com
This leaflet was last revised in March 2020.

This is a Medicament

- Medicament is a product which affects your health and its consumption contrary to instructions is dangerous for you.
- Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medicament.
- The doctor and the pharmacist are the experts in medicines, their benefits and risks.
- Do not by yourself interrupt the period of treatment prescribed for you.
- Do not repeat the same prescription without consulting your doctor.
- Keep all medicaments out of reach of children.

Council of Arab Health Ministers
Union of Arab Pharmacists



→ **Contact your doctor or pharmacist** if you have any questions about taking Avodart.

Other medicines and Avodart

Tell your doctor if you are taking, have recently taken, or might take any other medicines.

Some medicines can react with Avodart and may make it more likely that you'll have side-effects. These medicines include:

- verapamil or diltiazem** (for high blood pressure)
- ritonavir or indinavir** (for HIV)
- itraconazole or ketoconazole** (for fungal infections)
- nefazodone** (an antidepressant)
- alpha-blockers** (for enlarged prostate or high blood pressure).

→ **Tell your doctor** if you are taking any of these medicines. Your dose of Avodart may need to be reduced.

Avodart with food and drink

Avodart can be taken with or without food.

Pregnancy, breast-feeding and fertility
Women who are pregnant (or may be) must not handle leaking capsules. Dutasteride is absorbed through the skin and can affect the normal development of a male baby. This is a particular risk in the first 16 weeks of pregnancy.

Use a condom during sexual intercourse. Dutasteride has been found in the semen of men taking Avodart. If your partner is or may be pregnant, you must avoid exposing her to your semen.

Avodart has been shown to reduce sperm count, semen volume and sperm movement. Therefore male fertility may be reduced.

→ **Contact your doctor** for advice if a pregnant woman has come into contact with dutasteride.

Driving and using machines
Avodart is unlikely to affect your ability to drive or operate machinery.

Avodart contains lecithin from soya

This medicine contains lecithin from soya, which may contain soya oil. If you are allergic to peanut or soya, do not use this medicinal product.

3 How to take Avodart

Always take Avodart exactly as your doctor or pharmacist has told you to. If you do not take it regularly the monitoring of your PSA levels may be affected. Check with your doctor or pharmacist if you are not sure.

How much to take

The recommended dose is one capsule (0.5 mg) taken once a day. Swallow the capsules whole with water. Do not chew or break open the capsule. Contact with the contents of the capsules may make your mouth or throat sore.

Avodart is a long-term treatment. Some men notice an early improvement in their symptoms. However, others may need to take Avodart for 6 months or more before it begins to have an effect. Keep taking Avodart for as long as your doctor tells you.

If you take more Avodart than you should
Contact your doctor or pharmacist for advice if you take too many Avodart capsules.

If you forget to take Avodart

Do not take a double dose to make up for a forgotten dose. Just take your next dose at the usual time.

Don't stop Avodart without advice

Don't stop taking Avodart without talking to your doctor first. It may take up to 6 months or more for you to notice an effect.

→ **If you have any further questions on the use of this medicine, ask your doctor or pharmacist.**

4 Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Allergic reaction

The signs of allergic reactions can include:

- skin rash** (which can be itchy)
- hives** (like a nettle rash)
- swelling of the eyelids, face, lips, arms or legs.**

→ **Contact your doctor immediately** if you get any of these symptoms, and **stop using Avodart.**

Common side effects

These may affect up to 1 in 10 men taking Avodart:

- not able to achieve or maintain an erection (*impotence*), this may continue after you stop taking Avodart
- decreased sex drive (*libido*), this may continue after you stop taking Avodart
- difficulty with ejaculation, such as a decrease in the amount of semen released during sex, this may continue after you stop taking Avodart

Package Leaflet: Information for the user

Avodart

0.5 mg soft capsules
Dutasteride

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

- What Avodart is and what it is used for
- What you need to know before you take Avodart
- How to take Avodart
- Possible side effects
- How to store Avodart
- Contents of the pack and other information

1 What Avodart is and what it is used for

Avodart is used to treat men with an enlarged prostate (benign prostatic hyperplasia) - a non-cancerous growth of the prostate gland, caused by producing too much of a hormone called dihydrotestosterone.

The active ingredient is dutasteride. It belongs to a group of medicines called 5-alpha reductase inhibitors.

As the prostate grows, it can lead to urinary problems, such as difficulty in passing urine and a need to go to the toilet frequently. It can also cause the flow of the urine to be slower and less forceful. If left untreated, there is a risk that your urine flow will be completely blocked (*acute urinary retention*). This requires immediate medical treatment. In some situations surgery is necessary to remove or reduce the size of the prostate gland. Avodart lowers the production of dihydrotestosterone, which helps to shrink the prostate and relieve the symptoms. This will reduce the risk of acute urinary retention and the need for surgery.

Avodart may also be used with another medicine called tamsulosin (used to treat the symptoms of an enlarged prostate).

2 What you need to know before you take Avodart

Do not take Avodart

- if you are allergic to **dutasteride, other 5-alpha reductase inhibitors, soya, peanut** or to any of the other ingredients of this medicine (listed in section 6)
- if you have a **severe liver disease.**

→ If you think any of these apply to you, **don't take** this medicine until you have checked with your doctor.

This medicine is for men only. It must not be taken by women, children or adolescents.

Warnings and Precautions

Talk to your doctor before taking Avodart.

- Make sure your doctor knows about liver problems.** If you have had any illness affecting your liver, you may need some additional check-ups while you are taking Avodart.
- Women, children and adolescents** must not handle leaking Avodart capsules, because the active ingredient can be absorbed through the skin. **Wash the affected area immediately** with soap and water if there is any contact with the skin.
- Use a condom during sexual intercourse.** Dutasteride has been found in the semen of men taking Avodart. If your partner is or may be pregnant, you must avoid exposing her to your semen as dutasteride may affect the normal development of a male baby. Dutasteride has been shown to decrease sperm count, semen volume and sperm motility. This could reduce your fertility.
- Avodart affects a blood test for PSA** (prostate-specific antigen), which is sometimes used to detect prostate cancer. Your doctor should be aware of this effect and can still use the test to detect prostate cancer. If you are having a blood test for PSA, tell your doctor that you are taking Avodart. **Men taking Avodart should have their PSA tested regularly.**
- In a clinical study of men at increased risk of prostate cancer, men taking Avodart had a **serious form of prostate cancer more often** than men who did not take Avodart. The effect of Avodart on this serious form of prostate cancer is not clear.
- Avodart may cause breast enlargement and tenderness.** If this becomes troublesome, or if you notice **breast lumps** or **nipple discharge** you should talk to your doctor about these changes as these may be signs of a serious condition, such as breast cancer.

Version: 1	
Harmony AMS Artwork Information Panel	
Manufacturing Site Number: 6200000081704	
Manufacturing Site(s): GSK POZNAN POLAND	
Product Market Trade Name: Avodart	
Approving Market(s): MARKET GROUP-Gulf and Near East	
Print Process: N/A	
Colour Standard Reference: N/A	
Technical Drawing (do NOT include version number): DRW L_023_03_FD_T	
Material Spec. (do NOT include version number): N/A	
Material Type:	N/A
Total Colours & Varnishes: 1	
BLACK	
Total Special Finishes: 0	
Body Text Size: 10.5pt	
Smallest Text Size: 10.5pt	
Loading: 12.5pt	
Horizontal Scale: 100%	
Microtext: N	
Additional Info (1): N/A	
Additional Info (2): N/A	
Additional Info (3): N/A	

APR_Production_V_NDD_06_2022 - Harmony - Version 3

200 mm Measuring Bar
If an e-banner DOES NOT appear on the top of this document, THEN this document has NOT been printed from the Harmony system.

IMPORTANT

GSK LOC is responsible to approve the change documentation, artwork brief and final artwork, ensuring that it is accurate, consistent and complete.

GSK SDC is responsible for site technical requirements and pre-press suitability.

GSK Market is responsible to advise SDC when changes required impact the following:

**Formulation
Tablet embossing
Storage conditions
Shelf Life**

NOTE TO MARKET

Local approvers must ensure that trade mark and copyright statements included in the brief comply with guidance provided by Legal: Global Trade Marks.

POZNAN

Site Additional Artwork Information Panel	
Production Site: GSK_POZNAN_POLAND	
Leaflet - full dimensions (width x height): 425 x 541 mm	folded dimensions (width x height): 45 x 77 mm
Foil / Laminate - dimensions (width): N/A	random repeats: N/A
Carton - dimensions (width x height x depth): N/A	
Tube - dimensions (Ø, length): N/A	Label - dimensions (width x height): N/A
Superseded Manufacturing Site Number: N/A	
Point of Sale Code Number: N/A	

6200000081704

Pharma code Ref. No. 1173 Pharma code Ref. No. 1173

