FRONT

128 Code - 000568

| Version: 2 | GSK |
|---|---|
| **Artwork Information Panel (AIP)** | |
| **Manufacturing Site Number:** 30000000000568 | |
| **Manufacturing Site(s):** Bora Pharmaceutical_Mississauga_CANADA | |
| **Product Market Trade Name:** Seroxat CR | |
| **Approving Market(s):** MARKET GROUP-Gulf and Near East | |
| **Print Process:** N/A | |
| **Colour Standard Reference:** N/A | |
| **Technical Drawing** (Do NOT include version number): 0002456 | |
| **Material Spec.** (Do NOT include version number): N/A | |
| **Material Type:** N/A | N/A |
| **Total Colours & Varnishes: 1** | |
| BLACK | |
| **Total Special Finishes: 0** | |
| **Body Text Size:** 7.5pt | |
| **Smallest Text Size:** 7.0pt | |
| **Leading:** 8.5pt | |
| **Horizontal Scale:** 100% | |
| **Microtext:** N | |
| **Additional Info (1):** N/A | |
| **Additional Info (2):** N/A | |
| **Additional Info (3):** N/A | |

200 mm Measuring Bar


**Artwork copyright is the property of the GSK Group of Companies.** All suppliers providing a service to GSK for printed components of any description must ensure that they have a licence for all fonts / software used in conjunction with GSK artwork. The distribution and use of fonts / software without a licence constitutes an intellectual property infringement. GSK will not accept any liability for the to/use of third party intellectual property rights by printed component suppliers. GSK reserves the right to recover any costs incurred as a result of third party intellectual property rights being infringed by printed component suppliers.


AIP_Production_V_INDD - 01_2023 - Version 4

## NOTE TO MARKET

**Local approvers must ensure that trade mark and copyright statements included in the brief comply with guidance provided by Legal: Global Trade Marks.**

## IMPORTANT

**GSK LOC is responsible to approve the change documentation, artwork brief and final artwork, ensuring that it is accurate, consistent and complete.**

**Artwork Studio is responsible for site technical requirements and pre-press suitability.**

**GSK Market is responsible to advise SDC when changes required impact the following:**

**Formulation, Tablet embossing, Storage conditions and Shelf Life**

A000568

SEROXAT CR
paroxetine hydrochloride

A000568

**SEROXAT CR**
**paroxetine hydrochloride**

128 Code - 000568

GSK

PATIENT MEDICATION INFORMATION

# SEROXAT CR

**12.5 mg & 25 mg controlled release film coated tablets**
**paroxetine hydrochloride**

**READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE**
**SEROXAT CR**
**Paroxetine Controlled Release Tablets**
Read this carefully before you start taking **SEROXAT CR** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **SEROXAT CR**.
**What is SEROXAT CR used for?**
SEROXAT CR is used in adults (18 years of age and older) to relieve symptoms of:
- **Major Depressive Disorder** (feeling sad, a change in appetite or weight, difficulty concentrating or sleeping, feeling tired, headaches, unexplained aches and pain).
- **Panic Disorder (with or without agoraphobia)** (panic attacks).
- **Social Phobia (social anxiety disorder)** (avoidance and/or fear of social situations).
- **Premenstrual Dysphoric Disorder (PMDD)** (episodes of major depression, severe mood changes, anxiety, irritability, physical pain, difficulty doing day to day tasks before your period).

SEROXAT CR is not for use in children under 18 years of age.
**How does SEROXAT CR work?**
SEROXAT CR belongs to the group of medicines called selective serotonin reuptake inhibitors (SSRIs). SEROXAT CR is thought to work by increasing the levels of a chemical in the brain called serotonin (5-hydroxytryptamine). This helps to relieve your symptoms of depression, panic disorder, social phobia or premenstrual dysphoric disorder. SEROXAT CR may take a number of weeks to work.
**What are the ingredients in SEROXAT CR?**
Medicinal ingredient: Paroxetine hydrochloride.
Non-medicinal ingredients: Colloidal silicon dioxide, glyceryl behenate, hydroxypropyl methylcellulose, lactose monohydrate, magnesium stearate, methacrylic acid copolymer type C, polyethylene glycol, polysorbate 80, polyvinylpyrrolidone, sodium lauryl sulphate, talc, titanium dioxide and triethyl citrate. In addition, each tablet also includes:
- 12.5 mg Tablet: D&C yellow No.10 aluminium lake, FD&C yellow No. 6 aluminium lake and yellow ferric oxide.
- 25 mg Tablet: D&C red No. 30 aluminium lake and red ferric oxide.

**SEROXAT CR comes in the following dosage forms:**
Controlled Release Tablets: 12.5 mg and 25 mg of paroxetine (as paroxetine hydrochloride).
**Do not use SEROXAT CR if:**
- you are allergic to paroxetine hydrochloride or to any of the non-medicinal ingredients in SEROXAT CR (see "What are the ingredients in SEROXAT CR").
- you are taking or have recently taken in the last 14 days medicines called monoamine oxidase inhibitors (MAOI) including linezolid (an antibiotic) or methylene blue (a dye injected into a vein during surgery, x-rays or other imaging procedures).
- you are taking or have recently taken thioridazine or pimozide. These medicines are used to treat mental health problems.

Ask your healthcare provider or pharmacist if you are not sure if you take a MAOI or one of these medicines, including the antibiotic linezolid or intravenous methylene blue. Do not start taking a MAOI or thioridazine or pimozide for at least 14 days after you stop treatment with SEROXAT CR.
**To help avoid side effects and ensure proper use, talk to your healthcare professional before you take SEROXAT CR. Talk about any health conditions or problems you may have, including if you:**
- have epilepsy or a history of seizures.
- have a history of liver or kidney problems.
- have heart problems.
- have a history or family history of mania/hypomania or bipolar disorder.
- have depression or other mental health disorders.
- have high cholesterol.
- have low levels of sodium in your blood.
- have had a recent bone fracture or were told you have osteoporosis or risk factors for osteoporosis.
- have a bleeding disorder or have been told that you have low platelets.
- are pregnant or planning to become pregnant.
- are breast feeding or planning to breast feed.
- have a history of alcohol or drug abuse.
- have ever had any allergic reactions to medications, food, etc.
- are taking oral contraceptives and are being prescribed SEROXAT CR for Premenstrual Dysphoric Disorder.
- are allergic to azo dye (FD&C Yellow No. 6 aluminium lake). The 12.5 mg tablet contains an azo dye component.
- have an eye condition known as narrow angles (the iris and cornea of the eye are closer than normal).

**Other warnings you should know about:**
**Pregnancy:** Only take SEROXAT CR during pregnancy if you and your healthcare professional have discussed the risks and have decided that you should. If you take SEROXAT CR near the end of your pregnancy, you are at a higher risk of heavy vaginal bleeding shortly after birth. If you become pregnant while taking SEROXAT CR, tell your healthcare professional **right away**.
**Effects on newborns:** In some cases, babies born to a mother taking SEROXAT CR during pregnancy may require hospitalization, breathing support and tube feeding. Be ready to seek medical help for your newborn if they:
- Have trouble breathing or feeding,
- Have muscle stiffness, or floppy muscles (like a rag doll),
- Have seizures (fits),
- Are shaking (jitteriness),
- Are constantly crying.

If you take SEROXAT CR:
- During early pregnancy, there is a possible slight increased risk that your newborn may have birth defects, particularly a heart defect.
- During late pregnancy, your newborn may be at risk of having a serious lung condition called Persistent Pulmonary Hypertension of the Newborn (PPHN), which causes breathing problems.

**Fertility and sexual function:** Taking medicines like SEROXAT CR may increase your risk of having sexual problems. This may continue after SEROXAT CR has been discontinued, including for months or years afterwards in some cases. Tell your healthcare professional if you experience symptoms such as a decrease in sexual desire, performance or satisfaction. Medicines like SEROXAT CR may affect sperm quality. Fertility in some men may be reduced while taking SEROXAT CR.
**Falls and fractures:** SEROXAT CR can cause you to feel dizzy or lightheaded and can affect your balance. This increases your risk of falling. In addition SEROXAT CR may increase your risk of breaking a bone if you are:
- elderly,
- have osteoporosis, or
- have other major risk factors for breaking a bone.

You should take extra care to avoid falls especially if you get dizzy or have low blood pressure.
**Driving and using machines:** SEROXAT CR may make you feel sleepy. Avoid driving a vehicle or using machinery until you know how it affects you.
**Angle-closure glaucoma:** SEROXAT CR can cause an acute attack of glaucoma. Having your eyes examined before you take SEROXAT CR could help identify if you are at risk of having angle-closure glaucoma. Seek immediate medical attention if you experience:
- eye pain,
- changes in vision,
- swelling or redness in or around the eye.

**Cholesterol and blood tests:** SEROXAT CR can cause abnormal blood test results, including elevated levels of cholesterol. Your healthcare professional will decide when to perform blood tests and will interpret the results.
**Do NOT stop taking SEROXAT CR without talking to your healthcare professional first. If stopped abruptly, SEROXAT CR may cause unwanted side effects such as:**
- light-headedness,
- nausea and vomiting,
- agitation/restlessness,
- anxiety,
- sweating,
- headache,
- sleep disturbance,
- electric shock sensations,
- tinnitus (buzzing, hissing, whistling, ringing or other persistent noise in the ears).

**Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.**

**Serious Drug Interactions**
**Do not take SEROXAT CR if you are taking or have recently taken any of the following drugs as you may have serious side effects:**
- monoamine oxidase inhibitors (MAOIs) such as the antibiotic linezolid and the intravenous dye methylene blue.
- thioridazine (typically used to treat schizophrenia and psychosis).
- pimozide (typically used to manage Tourette's syndrome).

Wait **14 days** after you stop taking a MAOI, or thioridazine, or pimozide before starting SEROXAT CR. If you are unsure, ask your healthcare professional.

**The following may also interact with SEROXAT CR.**
- other antidepressants, such as SSRIs, SNRIs, and certain tricyclics.
- other drugs that affect serotonin such as, lithium (used to treat bipolar depression), linezolid (antibiotic), tramadol (used to treat pain), tryptophan (used to treat anxiety or used as a sleep aid), and triptans (used to treat migraines).
- drugs used to prevent fits or treat epilepsy (anticonvulsants), such as carbamazepine, phenytoin, sodium valproate.
- drugs used to treat pain, such as fentanyl (used in anaesthesia or to treat chronic pain), tramadol, tapentadol, meperidine, methadone, and pentazocine.
- drugs used to treat breast cancer or fertility problems, such as tamoxifen.
- drugs used to treat patients with irregular heart beats (arrhythmias).
- drugs used to treat schizophrenia.
- drugs used to treat Human Immunodeficiency Virus (HIV) infection, such as a combination of fosamprenavir and ritonavir.
- drugs used to treat Parkinson's Disease or other movement disorders, such as procyclidine.
- drugs used to treat high blood pressure and angina, such as metoprolol.
- drugs which may affect blood clotting and increase bleeding, such as oral anti-coagulants (e.g. warfarin, dabigatran), acetylsalicylic acid (e.g. aspirin) and other non-steroidal anti- inflammatory drugs (e.g. ibuprofen).
- drugs that affect the central nervous system, such as haloperidol, amylbarbitone, and oxazepam.
- drugs used to treat cough, such as dextromethorphan.
- drugs to treat heartburn, such as cimetidine.
- drugs to treat respiratory diseases (chronic obstructive pulmonary disease (COPD) and asthma), such as theophylline.
- any natural herbal products (e.g. St. John's Wort).
- alcohol.

**How to take SEROXAT CR:**
- It is very important that you take SEROXAT CR exactly as your healthcare professional has instructed.
- Take your tablets in the morning, with or without food.
- Swallow tablet(s) whole with water. Do not chew or crush tablet(s).
- You should continue to take your medicine even if you do not feel better, as it may take a number of weeks for your medicine to work.
- Keep taking your tablets, as instructed, until your healthcare professional tells you to stop.
- Talk to your healthcare professional before you stop taking your medication on your own.

**Remember:** This medicine has been prescribed only for you. Do not give it to anybody else, as they may experience undesirable effects, which may be serious.
**Usual Dose:**
The starting dose of SEROXAT CR depends on your illness and current health. It is usually 12.5 mg or 25 mg once a day in the morning. Your healthcare professional may gradually increase your dose to help control your symptoms.
**Overdose:**

If you think you, or a person you are caring for, have taken too much SEROXAT CR, contact a healthcare professional, hospital emergency department, or regional poison control centre immediately, even if there are no symptoms.

**Missed Dose:**
If you forget to take your tablet in the morning, take it as soon as possible, unless it is too close to the time of the next dose. Take your next dose at the normal time the next morning, then carry on as before. Do not try to make up for a missed dose by taking a double dose the next time.
**What are possible side effects from using SEROXAT CR?**
These are not all the possible side effects you may have when taking SEROXAT CR. If you experience any side effects not listed here, tell your healthcare professional.
Side effects may include:
- blurred vision
- constipation
- diarrhea
- dizziness
- drowsiness
- dry mouth
- feeling agitated
- headache
- loss of appetite
- nausea/vomiting
- nervousness
- sexual problems (decreases in sexual desire, performance and satisfaction, may also lead to further decreases, which may continue after the drug is stopped)
- skin rash or hives alone
- sleep disturbances (abnormal dreams including nightmares)
- sweating
- tremor (shaking)
- weakness
- weight gain.

| Serious side effects and what to do about them | | | |
|---|---|---|---|
| Symptom / effect | Talk to your healthcare professional | | Stop taking drug and get immediate medical help |
| | Only if severe | In all cases | |
| **UNCOMMON** | | | |
| Dilated pupils | | ✓ | |
| **Gastrointestinal bleeding** (bleeding in the stomach or bowels): vomiting blood or passing black, tarry stool, blood in stool | | | ✓ |
| **Hallucinations:** seeing and hearing things that are really not there | | ✓ | |
| **Hypotension** (low blood pressure): dizziness, light-headedness or fainting when standing up from a sitting down or lying position | | ✓ | |
| **Oedema**: swelling of hands, ankles or feet | | ✓ | |
| **Seizures** (fits): uncontrollable shaking with or without loss of consciousness | | | ✓ |
| **Urinary retention** (inability to urinate or empty or loss of control of the bladder): pain | | ✓ | |
| **RARE** | | | |
| **Thrombocytopenia** (low platelets): bruising or unusual bleeding from the skin or other areas, bleeding for longer than usual if you hurt yourself, fatigue and weakness | | ✓ | |
| **Urinary incontinence** (involuntary loss of urine) | | ✓ | |
| **VERY RARE** | | | |
| **Severe allergic reactions:** red and lumpy skin rash, itching, hives, swelling of the lips, face, tongue, throat, trouble breathing, wheezing, shortness of breath, skin rashes, collapse or loss of consciousness | | | ✓ |
| **Severe skin reactions (Stevens- Johnson syndrome, toxic epidermal necrolysis, and erythema multiforme):** any combination of itchy skin rash, redness, blistering and peeling of the skin and/or inside of the lips, eyes, mouth, nasal passages or genitals, accompanied by fever, chills, headache, cough, body aches or swollen glands, joint pain, yellowing of the skin or eyes, dark urine. | | | ✓ |
| **UNKNOWN FREQUENCY** | | | |
| **Akathisia:** (a type of movement disorder): feeling restless, unable to sit or stand still. | | ✓ | |
| **Angle-closure glaucoma** (eye condition that can cause damage to the optic nerve): increased pressure in your eyes, sudden eye pain, eye and head pain, swelling or redness in or around the eye, hazy or blurred vision, sudden loss of sight. | | | ✓ |
| **Changes in feelings or behaviour:** anger, anxiety or violent thoughts. | | ✓ | |
| **Hyponatremia** (low sodium level in blood): symptoms of tiredness, weakness, muscle twitching, confusion combined with achy, stiff or uncoordinated muscles. | | ✓ | |
| **Increase in the hormone prolactin:** In women: breast discomfort, leakage of milk from the breasts, missed periods, or other problems with your menstrual cycle. In men: decreased body and facial hair, breast swelling, leakage of milk from the breasts, difficulty in getting or maintaining erections, or other sexual dysfunction. | | ✓ | |
| **Liver disorder**: symptoms include nausea, vomiting, loss of appetite combined with itching, yellowing of the skin or eyes, dark urine. | | ✓ | |
| **Mania:** elevated or irritable mood, decreased need for sleep, racing thoughts, overactive behaviour and thoughts. | | ✓ | |
| **Menstrual period disorders**: including heavy periods, bleeding between periods and absence of periods. | | ✓ | |
| **Photosensitivity** (sensitivity to sunlight): itchy, red skin when exposed to sunlight. | ✓ | | |
| **Restless legs syndrome:** irresistible urge to move the legs. | | ✓ | |

| Serious side effects and what to do about them | | | |
|---|---|---|---|
| Symptom / effect | Talk to your healthcare professional | | Stop taking drug and get immediate medical help |
| | Only if severe | In all cases | |
| **Serotonin toxicity (also known as serotonin syndrome) and neuroleptic malignant syndrome (NMS)**: a combination of most or all of the following: confusion, restlessness, sweating, shaking, shivering, high fever, hallucinations, sudden jerking of the muscles, muscle stiffness, feeling very agitated or irritable, fast heartbeat. The severity can increase, leading to loss of consciousness. | | | ✓ |
| Thoughts or actions about hurting or killing yourself. | | | ✓ |
| Uncontrollable movements of the body or face. | | ✓ | |

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, tell your healthcare professional.

**Reporting Side Effects**
If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet.
To report Product Complaint/s or Adverse Event/s associated with the use of GSK product/s, please contact us via:
gulf.safety@gsk.com. By reporting side effects you can help provide more information on the safety of this medicine.

**Storage:**
- Store between 15°C to 25 °C.
- Keep container tightly closed.
- If your healthcare professional tells you to stop taking SEROXAT CR, please return any leftover medicine to your pharmacist.
- You may need to read this package insert again. Please keep this package insert until you have finished your medicine.
- Keep out of reach and sight of children.

This leaflet was prepared by GlaxoSmithKline Inc.
Last Revised: September 22, 2022
**Manufactured by:** Bora Pharmaceutical Services Inc., 7333 Mississauga Road, Mississauga, Ontario, Canada, L5N 6L4
**Marketing Authorisation Holder:**
GlaxoSmithKline Inc, 100 Milverton Drive, 800, Mississauga, Ontario, Canada, L5R 4H1

*©2023 GSK group of companies or its licensor*
*Trademarks are owned by or licensed to the GSK group of companies*


This is a Medicament
- Medicament is a product which affects your health and its consumption contrary to instructions is dangerous for you.
- Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medicament.
- The doctor and the pharmacist are the experts in medicines, their benefits and risks.
- Do not by yourself interrupt the period of treatment prescribed for you.
- Do not repeat the same prescription without consulting your doctor.
- Keep all medicaments out of reach of children.

128 Code - 000568

| e-Banner | Project: CO-0051276 | Document: PPC-0030225 | Version: 4 |
|---|---|---|---|
| | Site Code: 30000000000568 | Operator: MZU43635 | Date/Time Created: 16.Aug.2023 15:02 GMT+1 |

| Version: 2 | GSK |
|---|---|
| **Artwork Information Panel (AIP)** | |
| **Manufacturing Site Number:** 30000000000568 | |
| **Manufacturing Site(s):** Bora Pharmaceutical_Mississauga_CANADA | |
| **Product Market Trade Name:** Seroxat CR | |
| **Approving Market(s):** MARKET GROUP-Gulf and Near East | |
| **Print Process:** N/A | |
| **Colour Standard Reference:** N/A | |
| **Technical Drawing** (Do NOT include version number): 0002456 | |
| **Material Spec.** (Do NOT include version number): N/A | |
| **Material Type:** N/A | N/A |
| **Total Colours & Varnishes: 1** | |
| BLACK | |
| **Total Special Finishes: 0** | |
| **Body Text Size:** 7.5pt | |
| **Smallest Text Size:** 7.0pt | |
| **Leading:** 8.5pt | |
| **Horizontal Scale:** 100% | |
| **Microtext:** N | |
| **Additional Info (1):** N/A | |
| **Additional Info (2):** N/A | |
| **Additional Info (3):** N/A | |

200 mm Measuring Bar

AIP_Production_V_INDD - 01_2023 - Version 4

Artwork copyright is the property of the GSK Group of Companies. All suppliers providing a service to GSK for printed components of any description must ensure that they have a licence for all fonts, software used in conjunction with GSK artwork. The distribution and use of fonts / software without a licence constitutes an intellectual property infringement. GSK will not accept any liability for the licence of third party intellectual property rights by printed component suppliers. Any fonts / software used in conjunction with GSK artwork must be licensed to the printer on a royalty free basis.

## NOTE TO MARKET

Local approvers must ensure that trade mark and copyright statements included in the brief comply with guidance provided by Legal: Global Trade Marks.

## IMPORTANT

GSK LOC is responsible to approve the change documentation, artwork brief and final artwork, ensuring that it is accurate, consistent and complete.

Artwork Studio is responsible for site technical requirements and pre-press suitability.

GSK Market is responsible to advise SDC when changes required impact the following:

**Formulation, Tablet embossing, Storage conditions and Shelf Life**

GSK

**المعلومات الدوائية للمريض**
**يجب قراءة هذه النشرة للحصول على استفادة أمن وفعال لدوائك.**

# سيروكسات سي أر

**أقراص باروكستين ذات الإطلاق المحكم.**

يرجى قراءة هذه النشرة بعناية قبل البدء في تناول **سيروكسات سي أر**، وفي كل مرة يصرف فيه لك. تُعد هذه النشرة الطبية ملخص ولن تخبرك بكل شيء عن هذا الدواء. لذلك يرجى التحدث إلى طبيبك حول حالتك الطبية وعلاجك، واسأل إذا كان هناك أي معلومات جديدة حول **سيروكسات سي أر**.

**ما هي دواعي استعمال سيروكسات سي أر؟**
يستخدم سيروكسات سي أر في البالغين (١٨ عاماً وما فوق) لتخفيف الأعراض الآتية:
- **اضطراب الاكتئاب الشديد** (الشعور بالحزن، وتغير في الشهية أو الوزن، وصعوبة التركيز أو النوم، والشعور بالتعب، والصداع، والارجاع والألم غير المبرر).
- **اضطراب الهلع (مع رهاب الخلاء أو بدونه)** (نوبات الهلع).
- **الرهاب الاجتماعي (اضطراب القلق الاجتماعي)** (التجنب و/أو الخوف من المواقف الاجتماعية).
- **اضطراب ما قبل الطمث الاكتئابي (PMDD)** (نوبات الاكتئاب الشديد، والتغيرات المزاجية الشديدة، والقلق، التهيج، والألم الجسدي، وصعوبة القيام بالمهام اليومية قبل الدورة الشهرية).

لا يستخدم سيروكسات سي أر في الأطفال دون سن ١٨ عاماً.

**كيف يعمل سيروكسات سي أر؟**
ينتمي سيروكسات سي أر إلى مجموعة الأدوية التي تسمى مثبطات استرداد السيروتونين الانتقائية (SSRIs). يعمل دواء سيروكسات سي أر عن طريق زيادة مستويات مادة كيميائية في المخ تسمى السيروتونين (٥-هيدروكسي تريبتامين). هذا يساعد في تخفيف أعراض الاكتئاب، أو اضطراب الهلع، أو الرهاب الاجتماعي أو اضطراب ما قبل الطمث الاكتئابي. يمكن أن يستغرق سيروكسات سي أر عدّة من الأسابيع حتى تظهر فاعليته.

**ما هي المكونات في سيروكسات سي أر؟**
المكون الدوائي: باروكستين هيدروكلوريد.
المكونات غير الدوائية:
ثنائي أكسيد السيليكون العروي، جليسيريل بيهابيتيت، هيدروكسي بروبيل ميثيل سيلليوز، مونوهيدرات الدكستروز، سليرات الماغنيسيوم، بولمير مشترك من حمض الميثاكريليك من النوع C، بولي إيثيلين جليكول، بولي سوربات ٨٠، بولي فينيل بيروليدون، كبريتات لوريل الصوديوم، تلك، ثاني أكسيد التيتانيوم، سيرات ثلاثي الإيثيل.
بالإضافة إلى ذلك، يحتوي كل قرص أيضًا على.
تحتوي أقراص سيروكسات سي أر بتركيز ١٢,٥ مجم أيضًا على صبغ الألومنيوم رقم ١٠ بالتون الأصفر للدواء ومستحضرات التجميل، وصبغ الألومنيوم رقم ٦ بالتون الأصفر للطعام ومستحضرات التجميل، وأكسيد الحديد الثلاثي الأصفر.
تحتوي أقراص سيروكسات سي أر بتركيز ٢٥ مجم أيضًا على صبغ الألومنيوم رقم ٣٠ بالتون الأحمر للدواء ومستحضرات التجميل، وأكسيد الحديد الثلاثي الأحمر.

**يوفّر دواء سيروكسات سي أر في أشكال الجرعات التالية:**
- الأقراص ذات الإطلاق المحكم بتركيز ١٢,٥ مجم، ٢٥ مجم، تحتوي على باروكستين (في شكل باروكستين هيدروكلوريد).

**موانع استعمال دواء سيروكسات سي أر إذا:**
- كنت تعاني من حساسية من مادة باروكستين هيدروكلوريد أو من أي من المكونات غير الفعلية الموجودة في دواء سيروكسات سي أر (يرجى الاطلاع على مكونات دواء سيروكسات سي أر).
- كنت تتناول حاليًا أو تناولت مؤخرًا في آخر ١٤ يومًا أي أدوية تسمى مثبطات الأوكسيداز أحادي الأمين (MAOI)(مثل ليزولايد (مضاد حيوي)، مسمة زرقاء من الميثيلين (صبغة تحقن في الوريد أثناء العمليات الجراحية أو الأشعة السينية أو إجراءات التصوير الأخرى.
- كنت تتناول حاليًا أو تناولت مؤخرًا أدوية أخرى لتستخدم لعلاج مشاكل الصحة العقلية مثل: ثيوريدازين أو بيموزيد.

**يوفّر دواء سيروكسات سي أر في أشكال الجرعات التالية:**
- الأقراص ذات الإطلاق المحكم بتركيز ١٢,٥ مجم، ٢٥ مجم، تحتوي على باروكستين (في شكل باروكستين هيدروكلوريد).

**موانع استعمال دواء سيروكسات سي أر إذا:**
- كنت مثبط لوكسيديز أحادي الأمين (على سبيل المثال، المضاد الحيوي لينزولايد، صبغة زرقاء من الميثيلين تحقن في الوريد)
- ثيوريدازين (يستخدم عادة لعلاج الفصام والذهان).
- بيموزيد (يستخدم عادة لإدارة متلازمة توريت).

انتظر ١٤ يوماً بعد التوقف عن تناول مثبط لوكسيديز أحادي الأمين أو ثيوريدازين أو بيموزيد قبل بدء سيروكسات سي أر. إذا كنت غير متأكد، اسأل الطبيب الرعاية الصحية المعالج.

**يمكن أن يتداخل ما يلي مع دواء سيروكسات سي أر.**
- مضادات الاكتئاب الأخرى، مثل مثبطات استرداد السيروتونين الانتقائية ومثبطات استرداد السيروتونين والنورأدرينالين الانتقائية وبعض مضادات الاكتئاب الثلاثية الحلقات.
- الأدوية الأخرى التي تؤثر على السيروتونين مثل الليثيوم (يستخدم لعلاج الاكتئاب ثنائي القطب)، ليفوتونين (مضاد حيوي)، ترامادول (يستخدم لعلاج الألم)، التريبتوفان (يستخدم لعلاج القلق أو يستخدم كمساعد على النوم)، التريبتان (يستخدم لعلاج الصداع النصفي).
- الأدوية المستخدمة لمنع النوبات أو علاج الصرع (مضادات التشنجات) مثل كاربامازيبين، فيناتوين، فالبروات الصوديوم.
- الأدوية المستخدمة لعلاج الألم، مثل الفنتانيل (يستخدم في التخدير أو لعلاج الآلام المزمنة)، والترامادول، والتابنادولين، والميبيريدين، والميثادون، والبنتازوسين.
- الأدوية المستخدمة في علاج سرطان الثدي أو مشاكل الخصوبة مثل دواء تاموكسيفين.
- الأدوية المستخدمة لعلاج المرضى الذين يعانون من عدم انتظام ضربات القلب (اضطراب نبض القلب).
- الأدوية المستخدمة في علاج القصور.
- الأدوية المستخدمة لعلاج عدوى فيروس نقص المناعة البشرية (HIV)، مثل مزيج من فوساموبرينافير وريتونافير.
- الأدوية المستخدمة لعلاج مرضى باركنسون أو اضطرابات الحركة الأخرى، مثل بروسيكليدين.
- الأدوية المستخدمة لعلاج ارتفاع ضغط الدم والذبحة الصدرية مثل ميتوبرولول.
- الأدوية التي قد تؤثر على تخثر الدم وتزيد من النزيف، مثل مضادات التخثر الفموية (مثل الوارفارين، ودابيجاتران)، جمض أسيتل السالسيليك (مثل الاسبرين) والأدوية الأخرى المضادة للالتهابات غير الستيرويدية (مثل الإيبوبروفين).
- الأدوية التي تؤثر على الجهاز العصبي المركزي، مثل هالوبيريدول، أميل باربيتون أو أوكسازيباي.
- الأدوية المستخدمة في علاج السعال، مثل ديكسترومثورفان.
- الأدوية المستخدمة لعلاج الحموضة المعوية مثل السيميتيدين.
- الأدوية المستخدمة لعلاج أمراض الجهاز التنفسي (مرض الانسداد الرئوي المزمن (COPD) والربو) مثل الثيوفيلين.
- أي منتجات عشبية طبيعية (مثل نبتة سانت جون).
- الكحول.

**كيفية تناول دواء سيروكسات سي أر:**
- من المهم جدًا أن تتناول دواء سيروكسات سي أر تمامًا كما أوصاك اخصائي الرعاية الصحية.
- تناول أقراصك في الصباح مع الطعام أو بدونه.
- ابلع القرص (الأقراص) كاملاً بالماء. لا تمضغ أو تسحق الأقراص.
- يجب أن تستمر في تناول دواءك حتى لو لم تشعر بتحسن، فقد يستغرق الدواء عدة أسابيع حتى تظهر فاعليته.
- يرجى الاستمرار في تناول أقراصك حسب التعليمات، حتى يخبرك اخصائي الرعاية الصحية المعالج بالتوقف.
- تحدث إلى اخصائي الرعاية الصحية قبل أن تتوقف عن تناول الدواء بمفردك.

**تنكر:** وُصف هذا الدواء من أجلك فقط. لا تعطيه لأي شخص آخر، فقد يتعرضون لآثار جانبية غير مرغوب فيها، والتي يمكن أن تكون خطيرة.

**الجرعة المعتادة:**
تعتمد جرعة الدواء من دواء سيروكسات سي أر على حالة مرضك وصحتك الحالية. عادة ما تكون ١٢,٥ مجم أو ٢٥ مجم بمعدل مرة واحدة في اليوم في الصباح. يمكن أن يزيد الطبيب المعالج جرعتك تدريجياً للمساعدة في السيطرة على الأعراض.

**الجرعة المفرطة:**
إذا كنت تعتقد أنك أو أي شخص تعتني به، قد تناولت الكثير من دواء سيروكسات سي أر، فيرجى الاتصال بأخصائي الرعاية الصحية أو قسم الطوارئ بالمستشفى أو المركز الإقليمي لمكافحة السموم على الفور، حتى إذا لم تظهر أعراض.

**السقوط والتصور:**
يمكن أن يسبب لك دواء سيروكسات سي أر الشعور بالدوخة أو الدوار ويمكن أن يؤثر على توازنك.
حيث يزيد هذا من خطر السقوط. بالإضافة إلى ذلك، يمكن أن يزيد دواء سيروكسات سي أر من خطر إصابتك بكسر في العظام إذا كنت:
- من كبار السن.
- تعاني من هشاشة العظام أو
- لديك عوامل خطر رئيسية أخرى تؤدي إلى كسر العظام.

يجب أن تنتبه جيدًا لتجنب السقوط خاصة إذا أصبت بالدوار أو كنت تعاني من انخفاض في ضغط الدم.

**القيادة واستخدام الآلات:**
يمكن أن يجعلك دواء سيروكسات سي أر تشعر بالنعاس. لذلك تجنب قيادة السيارة أو استخدام الآلات حتى تعرف كيف يؤثر الدواء عليك.

**جلوكوما الزاوية المغلقة:**
يمكن أن يسبب دواء سيروكسات سي أر نوبة حادة من الجلوكوما. يمكن أن يساعد فحص عينيك قبل تناول سيروكسات سي أر في تحديد ما إذا كنت معرضًا لخطر الإصابة بجلوكوما الزاوية المغلقة. يرجى طلب العناية الطبية الفورية إذا كنت تعاني من:
- ألم في العين.
- تغيرات في الرؤية.
- انتفاخ أو احمرار في العين أو حولها

**اختبارات الكوليسترول والدهون:**
يمكن أن يسبب سيروكسات سي أر تناتج غير طبيعية لفحص الدم، تشمل المستويات المرتفعة من الكوليسترول. سيقرر اخصائي الرعاية الصحية موعد إجراء اختبارات الدم وسيقوم بتفسير النتائج.

**لا تتوقف عن تناول سيروكسات سي أر بدون التحدث إلى اخصائي الرعاية الصحية المعالج أولاً. إذا توقفت عن تناول سيروكسات سي أر فجأة، يمكن أن يتسبب في آثار جانبية غير مرغوب فيها مثل:**
- دوار
- قيء، وغثيان.
- التهيج/ الارق.
- القلق.
- التعرق.
- الصداع
- اضطرابات النوم.
- الإحساس بصدمة كهربائية.
- طنين الأذن (أزيز، وهسهسة، وصفير، ورنين أو ضجيج آخر مستمر في الأذنين).

**يرجى إبلاغ اخصائي الرعاية الصحية المعالج عن جميع الأدوية التي تتناولها، تشمل أي أدوية أو فيتامينات أو معدّن أو مكملات طبيعية أو أدوية بديلة.**

**التفاعلات الدوائية الخطيرة**

**لا تستخدم دواء سيروكسات سي أر إذا كنت تتناول أو تناولت مؤخرًا: إذا من الأدوية التالية حيث يمكن أن تعاني من آثار جانبية خطيرة:**
- مثبط لوكسيديز أحادي الأمين (على سبيل المثال، المضاد الحيوي لينزولايد، صبغة زرقاء من الميثيلين تحقن في الوريد)
- ثيوريدازين (يستخدم عادة لعلاج الفصام والذهان).
- بيموزيد (يستخدم عادة لإدارة متلازمة توريت).

انتظر ١٤ يوماً بعد التوقف عن تناول مثبط لوكسيديز أحادي الأمين أو ثيوريدازين أو بيموزيد قبل بدء سيروكسات سي أر. إذا كنت غير متأكد، اسأل الطبيب الرعاية الصحية المعالج.

**يمكن أن يتداخل ما يلي مع دواء سيروكسات سي أر.**
- الأدوية الأخرى التي تؤثر على السيروتونين مثل الليثيوم (يستخدم لعلاج الاكتئاب ثنائي القطب)، ليفوتونين (مضاد حيوي)، ترامادول (يستخدم لعلاج الألم)، التريبتوفان (يستخدم لعلاج القلق أو يستخدم كمساعد على النوم)، التريبتان (يستخدم لعلاج الصداع النصفي).
- الأدوية المستخدمة لمنع النوبات أو علاج الصرع (مضادات التشنجات) مثل كاربامازيبين، فيناتوين، فالبروات الصوديوم.
- الأدوية المستخدمة لعلاج الألم، مثل الفنتانيل (يستخدم في التخدير أو لعلاج الآلام المزمنة)، والترامادول، والتابنادولين، والميبيريدين، والميثادون، والبنتازوسين.
- الأدوية المستخدمة في علاج سرطان الثدي أو مشاكل الخصوبة مثل دواء تاموكسيفين.
- الأدوية المستخدمة لعلاج المرضى الذين يعانون من عدم انتظام ضربات القلب (اضطراب نبض القلب).
- الأدوية المستخدمة في علاج القصور.
- الأدوية المستخدمة لعلاج عدوى فيروس نقص المناعة البشرية (HIV)، مثل مزيج من فوساموبرينافير وريتونافير.
- الأدوية المستخدمة لعلاج مرضى باركنسون أو اضطرابات الحركة الأخرى، مثل بروسيكليدين.
- الأدوية المستخدمة لعلاج ارتفاع ضغط الدم والذبحة الصدرية مثل ميتوبرولول.
- الأدوية التي قد تؤثر على تخثر الدم وتزيد من النزيف، مثل مضادات التخثر الفموية (مثل الوارفارين، ودابيجاتران)، جمض أسيتل السالسيليك (مثل الاسبرين) والأدوية الأخرى المضادة للالتهابات غير الستيرويدية (مثل الإيبوبروفين).
- الأدوية التي تؤثر على الجهاز العصبي المركزي، مثل هالوبيريدول، أميل باربيتون أو أوكسازيباي.
- الأدوية المستخدمة في علاج السعال، مثل ديكسترومثورفان.
- الأدوية المستخدمة لعلاج الحموضة المعوية مثل السيميتيدين.
- الأدوية المستخدمة لعلاج أمراض الجهاز التنفسي (مرض الانسداد الرئوي المزمن (COPD) والربو) مثل الثيوفيلين.
- أي منتجات عشبية طبيعية (مثل نبتة سانت جون).
- الكحول.

**كيفية تناول دواء سيروكسات سي أر:**
- من المهم جدًا أن تتناول دواء سيروكسات سي أر تمامًا كما أوصاك اخصائي الرعاية الصحية.
- تناول أقراصك في الصباح مع الطعام أو بدونه.
- ابلع القرص (الأقراص) كاملاً بالماء. لا تمضغ أو تسحق الأقراص.
- يجب أن تستمر في تناول دواءك حتى لو لم تشعر بتحسن، فقد يستغرق الدواء عدة أسابيع حتى تظهر فاعليته.
- يرجى الاستمرار في تناول أقراصك حسب التعليمات، حتى يخبرك اخصائي الرعاية الصحية المعالج بالتوقف.
- تحدث إلى اخصائي الرعاية الصحية قبل أن تتوقف عن تناول الدواء بمفردك.

**تنكر:** وُصف هذا الدواء من أجلك فقط. لا تعطيه لأي شخص آخر، فقد يتعرضون لآثار جانبية غير مرغوب فيها، والتي يمكن أن تكون خطيرة.

**الجرعة المعتادة:**
تعتمد جرعة الدواء من دواء سيروكسات سي أر على حالة مرضك وصحتك الحالية. عادة ما تكون ١٢,٥ مجم أو ٢٥ مجم بمعدل مرة واحدة في اليوم في الصباح. يمكن أن يزيد الطبيب المعالج جرعتك تدريجياً للمساعدة في السيطرة على الأعراض.

**الجرعة المفرطة:**
إذا كنت تعتقد أنك أو أي شخص تعتني به، قد تناولت الكثير من دواء سيروكسات سي أر، فيرجى الاتصال بأخصائي الرعاية الصحية أو قسم الطوارئ بالمستشفى أو المركز الإقليمي لمكافحة السموم على الفور، حتى إذا لم تظهر أعراض.

**الجرعة الفائتة:**
إذا نسيت تناول قرصك في الصباح، فتناوله في أسرع وقت ممكن، إلا إذا كان قريبًا جدًا من وقت الجرعة التالية. فتناول جرعتك التالية في الوقت المعتاد في صباح اليوم التالي، ثم استمر كما كان من قبل. لا تحاول تعويض الجرعة الفائتة عن طريق أخذ جرعة مضاعفة في المرة القادمة.

**ما هي الآثار الجانبية المحتملة عن استخدام دواء سيروكسات سي أر؟**
هذه ليست كل الآثار الجانبية المحتملة التي يمكن أن تصيبك عند تناول دواء سيروكسات سي أر. إذا تعرضت لأي آثار جانبية غير مذكورة هنا، فيرجى إبلاغ اخصائي الرعاية الصحية المعالج.
قد تشمل الآثار الجانبية:
- عدم وضوح الرؤية.
- الإمساك.
- الإسهال.
- الإرهاق.
- جفاف الفم.
- التعرق بالاضطراب.
- الصداع
- فقدان الشهية.
- قلق، أرق.
- التوتر العصبي.
- المشاكل الجنسية (انخفاض الرغبة، والقدرة، والإشباع الجنسي، يمكن أن يؤدي إحدى إلى مزيد من الانخفاضات، والتي قد تستمر بعد التوقف عن تناول الدواء).
- ملمح جلدي أو الشرى بمفرده.
- اضطرابات النوم (الحلم غير طبيعية تشمل الكوابيس).
- التعرق.
- رعاش (اهتزاز).
- ضعف.
- زيادة الوزن.

| الآثار الجانبية الخطيرة، وكيفية التعامل معها | | |
|---|---|---|
| العرض/ التأثير | تحدث إلى أخصائي الرعاية الصحية المعالج | |
| | إذا كانت الحالة شديدة فقط | في جميع الحالات |
| **حالات غير شائعة** | | |
| انتباه حدقة العين. | | ✓ |
| **نزيف الجهاز الهضمي:** (نزيف في المعدة أو الأمعاء): قيء دموي أو خروج براز أسود قطراني، مع وجود دم في البراز | | ✓ |
| **الهلوسة:** رؤية وسماع أشياء غير موجودة بالفعل | | ✓ |
| **انخفاض ضغط الدم** (ضغط الدم المنخفض): الدوخة، أو الدوار، أو الإحساس عند الوقوف من وضعية الجلوس أو الاستلقاء | | ✓ |
| **النوبة:** تورم في اللبنين أو الكاحلين أو القدمين | | ✓ |
| **الثوبيك:** اهتزاز لا يمكن السيطرة عليه مع فقدان الوعي أو بدونه | | ✓ |
| **احتباس البول** (عدم القدرة على التبول أو التفريغ أو فقدان السيطرة على المثانة): الم | | ✓ |
| **حالات نادرة** | | |
| **قلة الصفيحات** (انخفاض الصفائح الدموية): كدمات أو نزيف غير عادي من الجلد أو مداخل أخرى، والنزيف لفترة أطول من المعتاد إذا كنت تؤذي نفسك، والتعب والضعف. | | ✓ |
| **سلس البول** (فقدان التحكم في البول بشكل لاإرادي) | | ✓ |
| **حالات نادرة جدا** | | |
| **ربود القلق التحسسية الشديدة:** طفح جلدي أحمر ومتكتل، وحكة، وشري، وتورم الشفتين، والوجه، واللسان، والحلق، وصعوبة التنفس، والصفير، وضيق التنفس، وطفح جلدي، وهبوط، وفقدان الوعي | | ✓ |
| **ربود قبل جلبية شديدة (متلازمة ستيفنز-جونسون، تقشر الأدمة السامة الدموية البثورية التسممي، الحمامي متعدد الأشكال):** أي مزيج من الطفح الجلدي والحكة والاحمرار والقرح ونقشير الجلد داخل الشفتين أو العينين أو الفم أو الممرات الأنفية أو الأعضاء التناسلية أو كليهما، مصحوب بحمى، وقشعريرة، وصداع، وسعال، والام في الجسم، وتورم في الغدد، والام في المفاصل، وصفرار الجلد أو العينين، والبول الداكن | | ✓ |
| **تكرار غير معروف** | | |
| **تململ حركي:** (نوع من اضطرابات الحركة): الشعور عدم الراحة وعدم القدرة على الجلوس أو الوقوف دون حراك | | ✓ |
| **جلوكوما الزاوية المغلقة:** (حالة مرضية في العين يمكن أن تسبب ثقبًا في العصب البصري): زيادة الضغط في عينيك، والم مفاجئ في العين، والم في العين والرأس، وتورم أو احمرار في العين أو حولها، ضبابية الرؤية أو عدم وضوحها، وفقدان مفاجئ للبصر | | ✓ |
| **تغيرات في المشاعر أو السلوك:** العنف أو القلق أو الأفكار العنيفة | | ✓ |
| **نقص الصوديوم في الدم** (انخفاض مستوى الصوديوم في الدم): تظهر أعراض من التعب والضعف وارتخاش العضلات والاضطراب المصحوب بالم أو تهيس أو عدم تناسق العضلات | | ✓ |
| **زيادة هرمون البرولاكتين:**<br>في النساء: عدم الراحة في الثدي، أو تسرب الحليب من الثدي، أو غياب الدورة الشهرية، أو مشاكل أخرى في الدورة الشهرية.<br>في الرجال: انخفاض شعر الجسم والوجه، تورم الثدي، تسرب الحليب من الثدي، صعوبة في الانتصاب أو الحفاظ عليه، أو اختلال وظيفة جنسية أخرى. | | ✓ |

| الآثار الجانبية الخطيرة، وكيفية التعامل معها | | |
|---|---|---|
| العرض/ التأثير | تحدث إلى أخصائي الرعاية الصحية المعالج | توقف عن تناول الدواء واحصل على مساعدة طبية فورية |
| | إذا كانت الحالة شديدة فقط | في جميع الحالات |
| **اضطراب وظائف الكبد:** تشمل الأعراض الغثيان والقيء وفقدان الشهية المصحوب بالحكة واصفرار الجلد أو العينين والبول الداكن. | | ✓ |
| **الهوس:** ارتفاع المزاج أو الاندفاع، والخفاض الحاجة إلى النوم، وتسابق الأفكار، والسلوك والأفكار مفرطة النشاط | | ✓ |
| **اضطرابات الدورة الشهرية:** تتراوح بين غزارة، أو نزيف أو غياب الدورة الشهرية | | ✓ |
| **الحساسية للضوء** (حساسية لأشعة الشمس): حكة واحمرار في الجلد عند التعرض لأشعة الشمس | ✓ | |
| **متلازمة تعامل السلفين:** رعبة ملحة خارجة عن السيطرة في تحريك الساقين | ✓ | |
| **سمية السيروتونين (المعروف أيضًا باسم متلازمة السيروتونين) والمتلازمة الخبيثة لدواء المضاد للذهان:** مزيج من معظم أو كل ما يلي: الاضطراب، والارق، والتعرق، والاهتزاز، والرعشة، والحمى الشديدة، والهلوسة، والارتعاش الخاطئ للعضلات، وتيبس العضلات، والتعرق بأنهاج الشديد أو الاهمال، وسرعة ضربات القلب، يمكن أن تزداد شدتها، مما يؤدي إلى فقدان الوعي | | ✓ |
| أفكار أو أفعال حول إيذاء أو قتل نفسك | ✓ | |
| حركات الجسم أو الوجه التي لا يمكن السيطرة عليها | | ✓ |

إذا كان لديك أي عرض مزعج أو أثر جانبي غير مذكور هنا أو أصبحت حالتك سيئة بما يكفي لتؤثر على أنشطتك اليومية؛ يرجى إبلاغ أخصائي الرعاية الصحية المعالج.

**الإبلاغ عن الآثار الجانبية:**
للإبلاغ عن شكوى خاصة بالمنتج/ الآثار العكسية المرتبطة باستخدام منتجات GSK يرجى التواصل معنا عبر: gulf.safety@gsk.com

**التخزين:**
- يحفظ في درجة حرارة تتراوح بين ٢٥-١٥ درجة مئوية.
- الحفاظ على العبوة مغلقة بإحكام.
- إذا أخبرك اخصائي الرعاية الصحية المعالج بالتوقف عن تناول دواء سيروكسات سي أر، فيرجى إعادة أي دواء متبقي إلى الصيدلي.
- يمكن أن تحتاج إلى قراءة هذه النشرة الداخلية مرة أخرى. يرجى الاحتفاظ بهذه النشرة الداخلية حتى تنتهي من تناول الدواء.
- يحفظ بعيدًا عن متناول الأطفال وأنظارهم.

أعدت هذه النشرة الطبية بواسطة شركة GlaxoSmithKline Inc.

تاريخ آخر مراجعة: ٢٢ سبتمبر ٢٠٢٢

**الشركة المصنعة :** Bora Pharmaceutical Services Inc., ;
7333 Mississauga Road, Mississauga, Ontario, Canada, L5N 6L4

**المالكة لحق التسويق :** GlaxoSmithKline Inc
Milverton Drive, 800, Mississauga, Ontario, Canada, L5R 4H1
مجموعة شركات GSK 2023 أو المرخص لها.
العلامات التجارية مملوكة لمجموعة شركات GSK أو مرخصة لها

**ان هذا الدواء**
- الدواء مستحضر يؤثر على صحتك و استهلاكه خلافا للتعليمات يعرضك للخطر.
- اتبع بدقة وصفة الطبيب و طريقة الاستعمال المنصوص عليها و تعليمات الصيدلاني الذى صرفها لك.
- فالطبيب و الصيدلاني هما الخبيران بالدواء و بنفعه و ضرره.
- لا تقطع مدة العلاج المحددة لك من تلقاء نفسك.
- لا تكرر صرف الدواء بدون وصفة طبية.
- لا تترك الأدوية في متناول أيدي الأطفال.

مجلس وزراء الصحة العرب
و اتحاد الصيادلة العرب

A000568

128 Code - 000568

BACK