PATIENT LEAFLET

Keppra film-coated tablets

250 mg; 500 mg; 1000 mg

Levetiracetam

Read all of this leaflet carefully before you start taking this medicine.
- Keep this leaflet. You may need to read it again.
- If you have any questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you personally or your child. Don't pass it on to other people - it may harm them even if their symptoms seem to be the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

In this leaflet

1. What Keppra is and what it is used for
2. Before you take Keppra
3. How to take Keppra
4. Possible side effects
5. How to store Keppra
6. Further information
1. What KEPPRA is and what it is used for

Keppra contains the active ingredient levetiracetam. It belongs to antiepileptic medicines, which are used to treat fits (seizures) in epilepsy. On its own, Keppra is used to treat:
- partial onset seizure with or without secondary generalisation (the epilepsy form in which the fits initially affect only one side of the brain). This medicine can be started from 1 month of age and up to less than 16 years of age.
- myoclonic seizures (short, shock-like jerks of a muscle or group of muscles) in adults and adolescents from 12 years of age with juvenile myoclonic epilepsy.
- primary generalised tonic-clonic seizures (major fits, including loss of consciousness) in adults and adolescents from 12 years of age with idiopathic generalised epilepsy (the type of epilepsy that is thought to have a genetic cause).

2. Before you take KEPPRA

Don't take Keppra if:
- if you are **allergic** (hypersensitive) to levetiracetam, other pyrrolidone derivatives, or any other ingredients of Keppra (listed in Section 6)
→ if you think any of these apply to you, **don't take Keppra** until you have checked with your doctor.

Take special care with Keppra

Before you take Keppra your doctor needs to know:
- if you have **kidney problems or severe liver disease**, your doctor may need to adjust your dose of Keppra
- if you are **pregnant or breastfeeding**, think you may be pregnant or are planning to have a baby (*see Pregnancy and breast-feeding* later in section 2)
- if you have any **other medicines** (*see Other medicines and Keppra*)
- if you **are over 65**
- if you have a history or medical history of heart problems, for example disturbance in heart rhythm (visible on an electrocardiogram) or heart disease, or any other medicine that makes you prone to have disturbance in heart rhythm or unusual amount of salts in the body.

→ **Check with your doctor** if you think any of these may apply to you. Your doctor will decide whether Keppra is suitable for you.

Keppra film-coated tablets are not recommended for children under 6 years.

Keppra oral solution is the preferred formulation for use in infants and children **under the age of 6 years.**

Keppra is not indicated in children and adolescents below 16 years on its own (as monotherapy) (*see What Keppra is and what it is used for* in Section 1).

While you are taking Keppra

- If you notice any slowdown in the growth or unexpected puberty development of your child, **contact your doctor.**
- A small number of people being treated with anti-epileptics such as Keppra have had **thoughts of harming or killing themselves**. If at any time you have these thoughts, **immediately contact your doctor**.
- If you notice any unwanted aggressive behaviours, or if your family and friends notice important changes in mood or behaviour, immediately contact your doctor.
- Your seizures may rarely become worse or happen more often, mainly during the first month after starting the treatment or increase of the dose of Keppra. If you notice any of these symptoms while taking Keppra, immediately contact your doctor.

Consulting your doctor to look out for

Keppra can make some existing conditions worse, or cause serious side effects such as **severe allergic reactions, serious skin reactions, Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS), sudden decrease of kidney function, encephalopathy** (degenerative disease of the brain), **depression or suicidal thoughts**. You must look out for certain symptoms while you are taking Keppra to reduce the risk of any problems. See *Conditions you need to look out for* in Section 4.

Keppra can affect laboratory test results

Tell your doctor or pharmacist if you're taking any other medicines, if you've taken any recently, or if you start taking new ones. This includes medicines bought without a prescription.

Don't take **macrogol** (a drug used as **laxative**) for one hour before and one hour after taking Keppra as this may results in a loss of its effect.

You should tell your doctor if you are being treated for any of the following:
- methotrexate (used to treat certain types of **cancer**)

Some other medicines may affect how Keppra works, or make it more likely that you'll have side effects. Keppra can also affect how some other medicines work. These include:
- probenecid (used to treat **gout**)

→ **Tell your doctor or pharmacist** if you are taking any of these.

Pregnancy and breast-feeding

Keppra is not recommended for use during pregnancy.
- **Tell your doctor** if you are pregnant or planning to become pregnant.
- Use a **reliable method of contraception** while you're taking Keppra, to prevent pregnancy.
- **If you do become pregnant during treatment** with Keppra, tell your doctor.

Keppra can be used during pregnancy, only if after careful assessment it is considered necessary by your doctor.

You should not breastfeed while treated without discussing this with your doctor. A risk of birth defects for your unborn child cannot be completely excluded. Keppra has shown unwanted reproductive effects in animal studies.

Tell your doctor if you are breastfeeding during treatment with Keppra. The ingredients can pass into your breast milk. Talk to your doctor about this.

Driving and using machines

Keppra can make you feel drowsy or sleepy and have other side effects that may reduce you less alert. This is more likely at the beginning of treatment or after an increase in the dose.
→ Don't drive or use machines unless you are sure you're not affected.

3. How to take Keppra

How much to take

Always take Keppra exactly as your doctor has told you to. Check with your doctor or pharmacist if you're not sure.

Take the number of tablets/ the oral solution following your doctor's instructions.

Keppra must be taken **twice a day**, once in the morning and once in the evening, at about the same time each day.

Monotherapy

✦ Monotherapy in adults and adolescents (from 16 years of age)

Keppra film-coated tablets

When you will first start taking Keppra, your doctor will prescribe you a lower dose during first 2 weeks of treatment, before giving you the lowest general dose.

The usual starting dose of Keppra is **250 mg twice daily**. Your doctor will increase your dose to **500 mg, twice daily** after two weeks of treatment. Your doctor may decide to further increase your dose to a maximum of **1500 mg, twice daily** - depending on how you respond to the medicine.

Keppra 100 mg/ml oral solution

Measure the appropriate dosage using the syringe included in the package.

→ Ask your doctor or pharmacist if you have any questions on how to measure your dose accurately.

The usual starting dose of Keppra is 5 ml (500mg) twice daily.

General dose: Keppra is taken **twice daily**, in two equally divided doses, each individual dose being measured between **5 ml (500mg) and 15 ml (1500mg)**.

✦ Add-on therapy in children 6 months and older

Keppra film-coated tablets

The usual starting dose of Keppra is **500 mg twice daily**. Your doctor may decide to gradually increase your dose to a maximum of **1500 mg, twice daily** - depending on how you respond to the medicine.

Keppra 100 mg/ml oral solution.

Measure the appropriate dosage using the syringe included in the package.

→ Ask your doctor or pharmacist if you have any questions on how to measure your dose accurately.

The usual starting dose of Keppra is 5 ml (500mg) twice daily.

General dose: Keppra is taken **twice daily**, in two equally divided doses, each individual dose being measured between **5 ml (500mg) and 15 ml (1500mg)**.

✦ Add-on therapy in children 6 months and older

Keppra oral solution can be administered for use in infants and children **under the age of 6 years**.

Measure the appropriate dosage using the syringe size as advised by your doctor.

→ Ask your doctor or pharmacist if you have any questions on how to measure the dose accurately.

General dose: Keppra is taken twice daily, in two equally divided doses, each individual dose being measured between 0.07 ml (7mg) and 0.21 ml (21mg), per kg bodyweight of the infant. (see table below for dose examples).

Dose in infants (1 month to less than 6 months):

Weight	Starting dose: 0.07 ml/kg twice daily	Maximum dose: 0.21 ml/kg twice daily
4 kg	0.3 ml twice daily	0.85 ml twice daily
5 kg	0.35 ml twice daily	1.05 ml twice daily
6 kg	0.45 ml twice daily	1.25 ml twice daily
7 kg	0.5 ml twice daily	1.5 ml twice daily

Patients with kidney problems

Your doctor will decide on the correct dose of Keppra for you/your child depending on kidney function and the body weight.

How to take

Film-coated tablets

Swallow Keppra tablet(s) with a sufficient quantity of liquid (for example a glass of water). You can take Keppra with or without food. After oral solution is taken the bitter taste of levetiracetam may be experienced.

Oral solution

After measuring the correct dose with an appropriate syringe, Keppra oral solution may be diluted in a glass of water or baby's bottle and can be taken with or without food. After oral administration the bitter taste of levetiracetam may be experienced.

Take the oral solution following your doctor's instructions.

A dosing of 300 ml amber glass bottle 10 ml graduated oral syringe is supplied with the pack so you can measure your dose accurately.

If you forget to take Keppra

Don't take an extra dose to make up for a missed dose. Contact your doctor if you have missed one or more doses.

If you take too much Keppra

If you take more Keppra than you should you may be **more likely to feel drowsy, agitated or have other side effects such as decrease of alertness, aggression, breathing problems and loss of consciousness (coma).**

→ **Don't delay. Contact your doctor or your nearest hospital emergency department immediately.** If possible, show them the Keppra pack.

If you stop Keppra without advice

Don't stop taking Keppra or change your dose or recommended. Don't stop unless your doctor advises you to. If you are suffering from epilepsy abruptly stopping your medicine may increase your fits (seizures).

If stopping treatment is required, it should be discontinued gradually. Your doctor will instruct you about the gradual withdrawal of Keppra.

→ Ask your doctor or pharmacist if you have any questions on the use of this product.

4. Possible side effects

Like all medicines, Keppra can cause side effects, but not everybody gets them.

Conditions you need to look out for

Severe allergic reactions. These are rare in people taking Keppra. Signs include:
- raised and itchy rash (hives)
- swelling, sometimes of the face or mouth (*angioedema*), causing difficulty in breathing
- collapse or loss of consciousness

Serious skin reactions. These are rare in people taking Keppra. Signs include:
- skin rash, which may blister, and looks like small targets (central dark spots surrounded by a paler area, with a dark ring around the edge - *erythema multiformi*)
- a widespread rash with blisters and peeling skin, particularly around the mouth, nose, eyes and genitals (*Stevens Johnson Syndrome*)
- *extensive peeling of the skin on much of the body surface – (toxic epidermal necrolysis)*

Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS). These are rare in people taking Keppra. Signs include:
- flu-like symptoms and a rash on the face followed by an extended rash with a high temperature
- enlarged lymph nodes
- increased levels of their enzymes seen in blood tests and an increase in a type of white blood cell (*eosinophilia*)

Sudden decrease of kidney function. This is rare in people taking Keppra. Signs include:
- low urine volume
- tiredness, nausea, vomiting
- confusion
- swelling in the legs, ankles or feet

Encephalopathy (degenerative disease of the brain). This generally occur at the beginning of the treatment (few days to a few months) in people taking Keppra. Signs include:
- serious mental changes or signs of confusion
- feeling drowsy (somnolence)
- loss of memory (amnesia), memory impairment (*forgetfulness*)
- abnormal behaviour
- other neurological signs including involuntary or uncontrolled movements

Depression. This is common in people taking Keppra.

Suicidal thoughts. These are uncommon in people taking Keppra.

→ **Get medical help immediately** if you get these symptoms.

Very common side effects

These may affect **more than 1 in 10** people:
- inflammation of the nasopharynx (*nasopharyngitis*)
- feeling drowsy (somnolence)
- headache

Common side effects

These may affect **up to 1 in 10** people:
- loss of appetite (anorexia) - especially if you take another drug called topiramate
- depression, hostility or aggression, anxiety, difficulty in sleeping, nervousness or irritability
- fits (seizures), balance disorder, dizziness, abnormal drowsiness (lethargy), tremor
- spinning sensation (vertigo)
- cough
- stomach pain, diarrhoea, indigestion, vomiting, feeling sick (*nausea*)
- rash
- feeling weak or lack of energy

Uncommon side effects

These may affect **up to 1 in 100** people:
- decreased or increased weight
- suicide attempt and suicidal ideation, mental disorder, abnormal behaviour, seeing or hearing things that are not really there (hallucination), anger, confusion, panic attack, emotional instability/mood swings, agitation
- loss of memory, memory impairment (*forgetfulness*), abnormal coordination or loss of coordination of body movements, tingling or numbness of the hands or feet, disturbance in attention (loss of concentration)
- double vision, blurred vision
- hair loss, skin problems, eczema, itching
- muscular weakness, muscle pain
- injury

Uncommon side effects that may show up in blood tests:
- decrease in number of blood platelets - cells that help blood to clot (*thrombocytopenia*)
- decrease in the number of white blood cells (*leucopenia*)
- elevated/abnormal values in a liver function test

Rare side effects

These may affect **up to 1 in 1,000** people:
- infection
- allergic reactions (see 'Severe allergic reactions' earlier in section 4)
- drug-induced hypersensitivity reaction that includes fever, rash, and blood abnormalities (see Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS) earlier in Section 4)
- suicide, personality disorders Behavioural problems, abnormal thinking, severe confusion (delirium)
- uncontrollable muscle spasms affecting the eyes, head, neck and body, uncontrollable movements, hyperactivity (unusually overactive),
- encephalopathy (degenerative disease of the brain) (see 'encephalopathy' earlier in Section 4)
- seizures may become worse or happen more often
- abnormal results in liver function tests
- inflammation of the pancreas (*pancreatitis*)
- liver failure, inflammation of the liver
- erythema multiforme, Stevens Johnson Syndrome, toxic epidermal necrolysis (see 'Serious skin reactions' earlier in Section 4)
- acute kidney injury (see "Sudden decrease of kidney function" earlier in Section 4)
- rhabdomyolysis (breakdown of muscle tissue and associated creatine phosphokinase increase. Prevalence is significantly higher in Japanese patients when compared to non-Japanese patients
- limp or difficulty walking

Rare side effects that may show up in blood tests:
- decrease in number of all types of blood cells
- decrease in sodium in the blood

→ **Tell your doctor or pharmacist** if any of the side effects listed becomes **severe or troublesome**, or if you notice any side effects not listed in this leaflet

5. How to store KEPPRA

Keep out of the sight and reach of children.

Don't take Keppra after the expiry date shown on the pack.

Store below 30°C.

Keep out of the reach and sight of children.

Do not use after the expiry date stated on the container.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Further information

- What Keppra contains
- Keppra 250 mg film-coated tablets are blue, oblong, scored and debossed with the code ucb and 250 on one side.
- Keppra 500 mg film-coated tablets are yellow, oblong, scored and debossed with the code ucb and 500 on one side.
- Keppra 1000 mg film-coated tablets are white, oblong, scored and debossed with the code ucb and 1000 on one side.
- Keppra 100 mg/ml oral solution is a clear liquid.
- Keppra 100 mg/ml concentrate for solution for infusion is a clear, colourless, concentrate.

Keppra 250, 500 and 1000 mg film-coated tablets are packaged in aluminium/PVC blisters placed into cardboard boxes containing 100 film-coated tablets.

Keppra 100 mg/ml oral solution is packed in a 300 ml amber glass bottle (type III) with a white child resistant closure (polypropylene) in a cardboard box also containing a 10 ml graduated oral syringe (polypropylene, polyethylene) and an adaptor for the syringe (polyethylene). Keppra 100 mg/ml concentrate for solution for infusion is packed in 5 ml glass vials (type I) closed by a Teflon-faced grey chlorobutyl rubber stopper or an uncoated grey bromobutyl rubber stopper and sealed with an aluminium/polypropylene flip cap. Each carton contains 10 vials.

- Keppra Film Coated Tablets are Packed by:

Pharmalex, Belgium

Licensed by:

UCB Pharma Ireland Export Limited, UK

Trade marks are owned by or licensed to the GSK group of companies.

©2021 GSK group of companies or its licensor. All rights reserved.

NCDS Version Number: 11, Version Date: 27 January 2021

To report Product Complaint/s or Adverse Event/s associated with the use of GSK product/s, please contact us via: gulf.safety@gsk.com.

THIS IS A MEDICAMENT
- Medicament is a product which affects your health and its consumption contrary to instructions is dangerous for you.
- Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medicament.
- The doctor and the pharmacist are the medicament's experts, their benefits and risks.
- Do not by yourself interrupt the period of treatment prescribed.
- Do not repeat the same prescription without consulting your doctor.

Keep medicament out of the reach of children.

Council of Arab Health Ministers,

Union of Arab Pharmacists.

Rev. 11/2021 905297-C-1

DESTINATION LOCAL
DIMENSIONS 180 x 980 mm
COLOR BLACK

النشرة للمريض

كيبرا أقراص مغلفة

٢٥٠ مجم، ٥٠٠ مجم، ١٠٠٠ مجم

ليفيتيراسيتام

يرجى قراءة كافة محتويات هذه النشرة بعناية قبل البدء في استخدام هذا الدواء
- احتفظ بهذه النشرة. قد تحتاج إلى قراءتها لاحقًا.
- إذا كان لديك أي أسئلة أخرى، فيرجى استشارة طبيبك أو الصيدلي.
- تم وصف هذا الدواء لك فقط. لا تُعطِ الآخرين باستخدامه لأنه قد يضر بهم، حتى لو كانت أعراض المرض لديهم هي نفسها لديك.
- إذا أصبحت أي من الأعراض الجانبية خطيرة أو إذا عانيت من أي آثار جانبية غير مدرجة في هذه النشرة، فيرجى إبلاغ طبيبك أو الصيدلي.

محتويات هذه النشرة

١. ما هو كيبرا وفيم الاستخدام
٢. قبل تناول كيبرا
٣. كيفية استخدام كيبرا
٤. الآثار الجانبية المحتملة
٥. كيفية حفظ كيبرا
٦. معلومات إضافية

١. ما هو كيبرا وفيم الاستخدام

يحتوي كيبرا على المادة الفعالة ليفيتيراسيتام. ينتمي إلى الأدوية المضادة للصرع والتي تستخدم لعلاج نوبات الصرع.

يستخدم كيبرا وحده أو مع أدوية أخرى لعلاج:
- نوبات الصرع الجزئية مع أو بدون تعميم ثانوي (نوبات الصرع التي تؤثر في البداية على جانب واحد فقط من الدماغ). يمكن البدء في إعطاء هذا الدواء للأطفال من عمر شهر واحد وحتى أقل من ١٦ عامًا.
- نوبات الصرع الرعاشية (نوبات خاطفة وقصيرة تشبه الصدمة الكهربائية تحدث في عضلة أو مجموعة عضلات) لدى البالغين والمراهقين من عمر ١٢ عامًا فما فوق والذين تم تشخيصهم مؤخرًا بالصرع الرعاشي الطفولي.
- نوبات الصرع التوترية الرمعية الأولية (نوبات الصرع الكبرى، بما في ذلك فقدان الوعي) لدى البالغين والمراهقين من عمر ١٢ عامًا فما فوق والذين تم تشخيصهم مؤخرًا بالصرع المعمم مجهول السبب (النوع من الصرع الذي يُعتقد أن له سبب وراثي).

٢. قبل تناول كيبرا

لا تتناول كيبرا في الحالات التالية
- إذا كنت تعاني من الحساسية تجاه ليفيتيراسيتام أو مشتقات البيروليدون أو أي من المكونات الأخرى لهذا الدواء المدرجة في القسم (٦)
- إذا كنت تعاني من أي مما سبق، فلا تستخدم كيبرا حتى تتحدث مع طبيبك أولاً.

تحدث إلى طبيبك قبل تناول كيبرا

يجب على طبيبك معرفة ما يلي قبل تناولك كيبرا:
- **إذا كنت تعاني من مشاكل في الكلى أو مرض حاد في الكبد**، فقد يحتاج طبيبك إلى ضبط جرعتك من كيبرا
- **إذا كنتِ حاملاً أو مرضعة، تعتقدين أنكِ قد تكونين حاملاً أو تخططين لإنجاب طفل**، فيرجى استشارة الطبيب قبل تناول كيبرا. إذا كنتِ حاملاً، فلا يجب تناول كيبرا إلا إذا كان ذلك ضروريًا بعد تقييم دقيق من قبل طبيبك. لا تتوقفي عن تناول كيبرا دون استشارة طبيبك. يجب على النساء اللاتي يتناولن كيبرا عدم الرضاعة الطبيعية أثناء تناول كيبرا.
- **إذا كنت تتناول أدوية أخرى**. (انظر الفقرة بعنوان تناول أدوية أخرى تحت القسم ٣)
- **إذا كنت تبلغ من العمر ٦٥ عامًا فأكثر**، فقد تحتاج إلى تعديل جرعة كيبرا
- **إذا كنت تعاني من مشاكل في القلب**، مثل اضطراب نظم القلب (يظهر على تخطيط القلب) أو مرض القلب، أو أي دواء آخر يجعلك عرضة لاضطراب نظم القلب أو لاختلال في مستويات الأملاح في الجسم.

→ **تحدث إلى طبيبك** إذا كنت تعتقد أن أيًا من هذه الحالات تنطبق عليك. سيقرر طبيبك ما إذا كان كيبرا مناسبًا لك.

لا ينصح باستخدام أقراص كيبرا المغلفة للأطفال دون سن ٦ سنوات.

يُفضل استخدام محلول كيبرا الفموي للأطفال دون سن ٦ سنوات.

لا ينصح باستخدام كيبرا لعلاج الأطفال والمراهقين دون سن ١٦ عامًا بمفرده (انظر "**كيف يعمل كيبرا وفيم يستخدم**" في القسم ١).

أثناء تناول كيبرا
- إذا لاحظت أي تباطؤ في نمو طفلك أو ظهور نمو غير المتوقع لديه من سن البلوغ **اتصل بطبيبك**.
- **تفكر عدد قليل من الأشخاص الذين تم علاجهم بمضادات الصرع مثل كيبرا في إيذاء أنفسهم أو الانتحار**. إذا راودتك هذه الأفكار، فاتصل بطبيبك على الفور.
- إذا لاحظت أي سلوكيات عدوانية غير مرغوب فيها، أو إذا لاحظ أصدقاؤك وأفراد عائلتك تغيرات مهمة في المزاج أو السلوك، فاتصل بطبيبك على الفور.
- قد تزداد نوبات الصرع لديك سوءًا أو تحدث بشكل متكرر في حالات نادرة، خاصة خلال الشهر الأول بعد بدء العلاج أو زيادة جرعة كيبرا. إذا لاحظت أي من هذه الأعراض أثناء تناولك كيبرا، فاتصل بطبيبك على الفور.

تناول أدوية أخرى

يمكن أن يجعل كيبرا بعض الحالات الحالية أسوأ أو قد يتسبب في آثار جانبية خطيرة مثل **الحساسية المفرطة الشديدة أو تفاعلات عكسية في الجلد أو التفاعلات الناتجة عن فرط الحساسية للأدوية المصحوبة بأعراض جهازية (DRESS)** أو **انخفاض الكبرى أو انخفاض المفاجئ في وظائف الكلى أو اعتلال الدماغ** (مرض تنكسي في المخ) أو **الاكتئاب أو الأفكار الانتحارية**. يجب عليك التعرف على أعراض معينة أثناء تناولك كيبرا لتقليل خطر حدوث أي مشاكل. راجع "العلامات التي تستوجب الحذر في القسم ٤ المدرجة أدناه.

قد يؤثر كيبرا على نتائج الاختبارات المعملية

أخبر طبيبك أو الصيدلي إذا كنت تتناول أو تناولت مؤخرًا أي أدوية أخرى، أو إذا بدأت في تناول أدوية جديدة. ويشمل ذلك الأدوية المصروفة دون وصفة طبية.

لا تتناول **مكروجول** (دواء يستخدم **كملين**) لمدة ساعة واحدة قبل وبعد تناول كيبرا بفترة، لأن هذا قد يؤثر في فعاليته.

يجب أن تخبر طبيبك إذا كنت تتناول أيًا من الأدوية التالية:
- ميثوتركسيت (يستخدم لعلاج بعض أنواع **السرطان**)

بعض الأدوية الأخرى قد تؤثر على طريقة عمل كيبرا، أو تزيد من احتمال تعرضك لآثار جانبية. يمكن أن يتأثر كيبرا أيضًا على طريقة عمل بعض الأدوية الأخرى. وهي تشمل:
- بروبينيسيد (يستخدم لعلاج **النقرس**)

→ **أخبر طبيبك أو الصيدلي** إذا كنت تتناول أيًا من هذه الأدوية.

الحمل والرضاعة الطبيعية

لا ينصح باستخدام كيبرا أثناء فترة الحمل
- **أخبري طبيبك إن كنت حاملاً**، أو تخططين للحمل.
- استخدمي **طريقة موثوقة لمنع الحمل** أثناء تناولك كيبرا، وذلك لمنع حمل حدوث الحمل.
- إذا أصبحت حاملاً أثناء تناولك كيبرا، أخبري طبيبك.

يمكن استخدام كيبرا أثناء الحمل، فقط إذا اعتبر طبيبك أن ذلك ضروري بعد التقييم الدقيق.

يجب ألا تتوقف عن تناول الدواء من تلقاء نفسك. لا يمكننا أن نستبعد تمامًا خطر حدوث عيوب خلقية لطفلك الذي لم يولد بعد أظهرت تجربة على حيوانات التأثيرات الإنجابية غير المرغوب فيها.

يجب عليك إخبار طبيبك إذا كنت ترضعين طفلك أثناء تناولك كيبرا. وذلك لأن المكونات يمكن أن تنتقل إلى حليب الثدي. استشيري طبيبك حول هذا الموضوع.

القيادة وتشغيل الماكينات

يمكن أن يجعلك كيبرا تشعر بالنعاس أو **الخمول** أو قد يسبب أعراض جانبية أخرى قد تقلل من يقظتك. يكون هذا أكثر احتمالاً في بداية العلاج أو بعد زيادة الجرعة.

→ لا تقود السيارة أو تستخدم الآلات إلا إذا كنت متأكدًا من أن كيبرا لا يؤثر عليك.

٣. كيفية تناول كيبرا

كمية الدواء المتناولة

تناول كيبرا دائمًا كما أخبرك الطبيب أو الصيدلي بالضبط. استشر طبيبك أو الصيدلي إذا كنت غير متأكد.

تناول عدد الأقراص/ المحلول الفموي حسب تعليمات الطبيب.

يجب تناول كيبرا **مرتين في اليوم** مرة واحدة في الصباح ومرة واحدة في المساء، في نفس الوقت تقريبًا كل يوم.

العلاج الأحادي

✦ تناول هذا الدواء كعلاج أحادي للبالغين والمراهقين (من سن ١٦ سنة)

كيبرا أقراص مغلفة

عندما تبدأ تناول كيبرا لأول مرة سيصف لك طبيبك **جرعة منخفضة** خلال أول أسبوعين من العلاج قبل أن يصف لك فيما بعد أعلى جرعة عامة.

الجرعة البدائية المعتادة من كيبرا هي **٢٥٠ مجم مرتين يوميًا**. سيزيد طبيبك الجرعة إلى **٥٠٠ مجم مرتين يوميًا** بعد أسبوعين من العلاج.

قد يقرر طبيبك زيادة الجرعة إلى ١٥٠٠ مجم مرتين يوميًا - اعتمادًا على كيفية استجابتك للدواء.

محلول كيبرا الفموي ١٠٠ مجم/مل

قم بقياس الجرعة المناسبة باستخدام السرنجة الموجودة داخل العبوة.

→ استشر طبيبك أو الصيدلي إذا كان لديك أي أسئلة حول كيفية قياس الجرعة بدقة.

الجرعة البدائية المعتادة من كيبرا هي **٥ مل (٥٠٠ مجم) مرتين يوميًا**.

الجرعة العامة: يؤخذ كيبرا **مرتين يوميًا**، في جرعتين منقسمتين بالتساوي، حيث تتراوح كل جرعة فردية بين **٥ مل (٥٠٠ مجم) و١٥ مل (١٥٠٠ مجم)**.

✦ علاج إضافي للبالغين والمراهقين (من ١٢ عامًا) ومن يزنون ٥٠ كجم أو أكثر

كيبرا أقراص مغلفة

جرعة البدء المعتادة من كيبرا هي **٥٠٠ مجم مرتين يوميًا**. قد يقرر طبيبك زيادة الجرعة تدريجيًا إلى **١٥٠٠ ملغ كحد أقصى مرتين يوميًا** - اعتمادًا على كيفية استجابتك للعلاج.

محلول كيبرا الفموي

قم بقياس الجرعة المناسبة باستخدام السرنجة الموجودة داخل العبوة.

→ استشر طبيبك أو الصيدلي إذا كان لديك أي أسئلة حول كيفية قياس الجرعة بدقة.

الجرعة البدائية المعتادة من كيبرا هي **٥ مل (٥٠٠ مجم) مرتين يوميًا**.

الجرعة العامة: يؤخذ كيبرا **مرتين يوميًا**، في جرعتين منقسمتين بالتساوي، حيث تتراوح كل جرعة فردية بين **٥ مل (٥٠٠ مجم) و١٥ مل (١٥٠٠ مجم)**.

✦ علاج إضافي للأطفال من عمر شهر واحد إلى أقل من ٤ أشهر

يمكن استخدام محلول كيبرا الفموي للأطفال دون سن ٦ شهور.

قم بقياس الجرعة المناسبة باستخدام السرنجة الموجودة داخل العبوة.

→ استشر طبيبك أو الصيدلي إذا كان لديك أي أسئلة حول كيفية قياس الجرعة بدقة.

الجرعة العامة: يؤخذ كيبرا **مرتين يوميًا**، في جرعتين منقسمتين بالتساوي، حيث تتراوح كل جرعة فردية بين ٠.٧ مل (٧ مجم) و٢.١ مل (٢١ مجم) بالتناسب مع وزن الجسم الخاص بالرضيع (انظر الجدول أدناه).

الجرعة للأطفال في عمر ١ شهر إلى أقل من ٦ أشهر:

الوزن	جرعة البدء: ٠.٧ مل/كجم مرتين يوميًا	الجرعة القصوى: ٢.١ مل/كجم مرتين يوميًا
٤ كجم	٠.٣ مل مرتين يوميًا	٠.٨٥ مل مرتين يوميًا
٥ كجم	٠.٣٥ مل مرتين يوميًا	١.٠٥ مل مرتين يوميًا
٦ كجم	٠.٤٥ مل مرتين يوميًا	١.٢٥ مل مرتين يوميًا
٧ كجم	٠.٥ مل مرتين يوميًا	١.٥ مل مرتين يوميًا

المرضى الذين يعانون من مشكلة في الكلى

سيقرر طبيبك جرعة كيبرا الصحيحة والمناسبة لك أو لطفلك وفقًا لحالة وظائف الكلى وزن الجسم.

كيفية التناول

الأقراص المغلفة

ابلع قرص/ أقراص كيبرا مع كمية مناسبة من السوائل (كوب من الماء على سبيل المثال). يمكنك تناول كيبرا مع طعام أو بدونه. قد تذوق مادة ليفيتيراسيتام المرة في حلقك بعد تناول الدواء عن طريق الفم.

المحلول الفموي

بعد قياس الجرعة الصحيحة باستخدام السرنجة المناسبة، يمكن تخفيف محلول كيبرا الفموي في كوب من الماء أو زجاجة الطفل ويمكن تناوله مع أو بدون طعام. قد تذوق مادة ليفيتيراسيتام المرة في حلقك بعد تناول الدواء عن طريق الفم.

تناول المحلول الفموي باتباع تعليمات الطبيب.

تأتي العبوة مع زجاجة من الزجاج الكهرماني سعة ٣٠٠ مل، مقياس مدرج للجرعات الفموية سعة ١٠ مل، حتى تتمكن من قياس جرعتك بدقة.

إذا نسيت تناول كيبرا

لا تتناول جرعة إضافية لتعويض جرعة فاتتك. اتصل بطبيبك إذا فاتتك جرعة أو أكثر.

إذا تناولت جرعة أكبر من اللازم من كيبرا

إذا تناولت كمية من كيبرا أكثر مما ينبغي، فمن المرجح أن تشعر بالنعاس أو الخمول أو تضعر بعض الآثار الجانبية الأخرى مثل قلة اليقظة والعدوانية، **صعوبة في التنفس وفقدان الوعي (الغيبوبة).**

→ **لا تتوقف عن تناول كيبرا دون نصيحة الطبيب.** يجب عليك إخبار طبيبك إذا كنت تعاني من مرض الصرع في حال التوقف المفاجئ عن تناول الدواء قد يزيد من نوبات الصرع.

إذا توقفت عن تناول كيبرا دون نصيحة الطبيب

لا تتوقف عن تناول كيبرا أو تغير جرعتك أو تتوقف عن تناوله إذا كنت تعاني من مرض الصرع في حال التوقف المفاجئ عن تناول الدواء قد يزيد من نوبات الصرع.

إذا كنت بحاجة إلى التوقف عن تناول كيبرا، فسيتم إيقاف تناوله تدريجيًا. سيقدم لك طبيبك تعليمات حول كيفية التوقف التدريجي عن تناول كيبرا.

→ اسأل طبيبك أو الصيدلي إذا كان لديك أي أسئلة حول استخدام هذا المنتج.

٤. الآثار الجانبية المحتملة

مثل جميع الأدوية، يمكن أن يسبب كيبرا آثارًا جانبية ولكن لا يُشترط أن يصاب بها كل من هو هذا الدواء.

حالات يجب عليك الانتباه لها

الحساسية المفرطة وهذا يعني الحدوث لدى الأشخاص الذين يتناولون كيبرا. تشمل الأعراض ما يلي:
- تورم في الوجه أو الجفون أو الشفتين أو الحلق (والذي يمكن أن يسبب صعوبة في التنفس)
- طفح جلدي مثل الشرى (بقع حمراء مرتفعة ومثيرة للحكة)
- انخفاض ضغط الدم الذي يسبب الدوار أو الإغماء

الآثار الجلدية الخطيرة وهذا يعني الحدوث لدى الأشخاص الذين يتناولون كيبرا. تشمل الأعراض ما يلي:
- طفح جلدي يتحول إلى بثور، ويبدو مثل الأهداف (بقع داكنة في المنتصف محاطة بمنطقة فاتحة اللون وحلقة داكنة حولها - *شفي مُتعدّد الأشكال*)
- طفح جلدي واسع الانتشار مع بثور وتقشر الجلد، خاصة حول الفم والأنف والعينين والأعضاء التناسلية (*متلازمة ستيفنس- جونسون*)
- تقشر الجلد على نطاق واسع في معظم سطح الجسم (*انحلال البشرة السمي*)

التفاعل الدوائي مع فرط الإيوزينيات والأعراض الجهازية وهذا يتأثر حدوث لدى الأشخاص الذين يتناولون كيبرا. تشمل الأعراض ما يلي:
- أعراض تشبه أعراض الإنفلونزا مصحوبة بطفح جلدي على الوجه يليه امتداد للطفح الجلدي مع ارتفاع في درجة الحرارة
- تضخم الغدد الليمفاوية
- تغيرات في نتائج فحوصات الدم والتي تظهر زيادة في عدد خلايا الدم البيضاء المعينة (الإيوزينات)

انخفاض مفاجئ في وظائف الكلى وهذا يتأثر حدوث لدى الأشخاص الذين يتناولون كيبرا. تشمل الأعراض ما يلي:
- انخفاض في حجم البول
- الإرهاق والغثيان والقيء
- الارتباك
- تورم في الساقين أو الكاحلين أو القدمين

اعتلال الدماغ (مرض تنكسي في المخ) يحدث هذا بشكل عام في بداية العلاج (من بضعة أيام إلى بضعة أشهر) لدى الأشخاص الذين يتناولون كيبرا. تشمل الأعراض ما يلي:
- تغيرات عقلية كبيرة أو أعراض تدل على الحيرة والتشوش والارتباك
- فقدان الذاكرة أو النسيان
- ضعف في التركيز
- سلوك غير طبيعي
- مشاكل عصبية أخرى تشمل حركات لا إرادية أو الخارجة عن السيطرة

الاكتئاب يحدث هذا بشكل شائع لدى الأشخاص الذين يتناولون كيبرا.

الأفكار الانتحارية قد يحدث هذا لدى الأشخاص الذين يتناولون كيبرا.

→ **إذا ظهرت عليك أي من هذه الأعراض، فاطلب المساعدة الطبية على الفور.**

آثار جانبية شائعة جدًا

قد تؤثر على شخص واحد من كل ١٠ أشخاص
- التهاب البلعوم الأنفي (*التهاب البلعوم الأنفي*)
- النعاس (*النعاس*)
- الصداع

آثار جانبية شائعة

قد تؤثر على شخص واحد من كل ١٠ أشخاص
- فقدان الشهية (*فقدان الشهية*) - خاصة إذا كنت تتناول دواء آخر يسمى توبيراميت
- الاكتئاب العدواني أو العدائي، القلق، الصعوبة في النوم، العصبية أو حدة الطبع، وسرعة الانفعال
- نوبات الصرع (النوبات)، دوخة، تغير في توازنك، الشعور بالنعاس المفرط (الخمول)، رعشة
- دوار
- سعال
- ألم المعدة، الإسهال، عسر الهضم، الشعور بالغثيان (*الغثيان*)، القيء
- الطفح الجلدي
- الشعور بالضعف، أو الافتقار إلى الطاقة

آثار جانبية غير شائعة

قد تؤثر على شخص واحد من كل ١٠٠ شخص
- زيادة أو فق